

E. Casanovas*

LASCAR STRONG TYPES AND FORKING IN NIP THEORIES

Abstract. This is an updated and slightly expanded version of a tutorial given in the *Mini-Course in Model Theory*[†], Torino, February 9–11, 2011. Some parts were previously exposed in the Model Theory Seminar of Barcelona. The main goals were to clarify the relation of forking with some versions of splitting in NIP theories and to present the known results on G -compactness, including a full proof of a theorem of E. Hrushovski and A. Pillay on G -compactness of NIP theories over extension bases. The tutorial was given in parallel to a tutorial of H. Adler on forking and dividing over models in NTP_2 theories, now written in [2].

Sections 1 and 2 present basic well-known facts on strong types, G -compactness, simple theories, NIP theories and NTP_2 theories. In section 3 we discuss dividing, forking and some versions of splitting. No new facts are stated, but we give a systematic account of the implications between these notions under different assumptions. In sections 4 and 5 we develop the product of types based on the completeness of a set B over a subset A (meaning that all finitary types over A are realized in B). This approach is inspired in Lascar’s treatment of special types in [16] and might include some new observations. In section 6 we give a slightly simplified proof of Hrushovski-Pillay’s results from [10] on G -compactness of NIP theories over extension bases and on the equivalence of Lascar splitting and KP-splitting in NIP theories assuming ω -saturation over the subset.

The context is the standard in Model Theory. T is a complete theory with infinite models, L is its language and \mathfrak{C} is its monster model. Generally x is a tuple of variables and a a tuple of parameters. A set is small if its cardinality is smaller than the cardinality of the monster model. The lengths of tuples are possibly infinite but always small. For any sequence $(a_i : i < \alpha)$ and $i \leq \alpha$, $a_{<i} = (a_j : j < i)$. For any other issue concerning notation and terminology, consult [4].

1. Strong types and G -compactness

DEFINITION 1. Let A be a small set of parameters. A relation R on the monster model is *A -invariant* if it is invariant under the group $\text{Aut}(\mathfrak{C}/A)$ of all automorphisms of \mathfrak{C} pointwise fixing A , that is, $f(R) = R$ for all $f \in \text{Aut}(\mathfrak{C}/A)$. It is *type-definable over A* if R is the solution set $\pi(\mathfrak{C})$ of a set $\pi(x)$ of formulas over A . R is *definable over A* if $\pi(x)$ consists of a single formula $\varphi(x) \in L(A)$. Note that x might be an infinite tuple of variables, so we allow definable relations of infinite arity. If $A = \emptyset$ we say that R is *0-type-definable* or *0-definable*.

*Partially supported by grants MTM 2011-26840 of Spanish Ministerio de Economía y Competitividad and 2009SGR-187 of Catalan Government.

[†]Thanks are due to D. Zambella, organizer of the course.

Clearly, if R is definable over A , it is type-definable over A and this implies it is A -invariant. In general, these implications can not be reversed.

REMARK 1. 1. R is definable over A iff R and its complement are type-definable over A .

2. If R is type-definable over A and it is B -invariant, then it is type-definable over B .
3. R is A -invariant iff it is a union $R = \bigcup_{i \in I} R_i$ of relations R_i which are type-definable over A .

Proof. See Lemma 1.4 of [4]. □

DEFINITION 2. An equivalence relation E between tuples of the same length in the monster model is *bounded* if the number of its equivalence classes is small. It is called *finite* if this number is finite.

REMARK 2. Let $(E_i : i \in I)$ be a family of equivalence relations on \mathfrak{C}^α .

1. If all E_i are A -invariant, $\bigcap_{i \in I} E_i$ is A -invariant.
2. If all E_i are type-definable over A , $\bigcap_{i \in I} E_i$ is type-definable over A .
3. If all E_i are bounded, $\bigcap_{i \in I} E_i$ is bounded.

Proof. For item 3 see Remark 9.2 in [4]. □

DEFINITION 3. Let us fix a set of parameters A and an ordinal α .

1. The intersection of all bounded A -invariant equivalence relations on α -tuples is the smallest bounded A -invariant equivalence relation on these tuples. It is called the *Lascar equivalence relation over A* and it is denoted by \equiv_A^{Ls} . We say that the tuples a, b have the same *Lascar strong type over A* if $a \equiv_A^{\text{Ls}} b$.
2. The intersection of all bounded type-definable over A equivalence relations on α -tuples is the smallest bounded type-definable over A equivalence relation on α -tuples, it is called the *Kim-Pillay equivalence relation over A* and it is denoted by \equiv_A^{KP} . Two tuples a, b have the same *KP-type over A* if $a \equiv_A^{\text{KP}} b$.
3. The intersection of all finite A -definable equivalence relations on α -tuples is a bounded type-definable over A equivalence relation sometimes called the *Shelah equivalence relation* and denoted by $\equiv_A^{\text{suchthat}}$. We say that a, b have the same *strong type over A* if $a \equiv_A^{\text{suchthat}} b$.
4. As usual, we write $a \equiv_A b$ for equality of type over A : $\text{tp}(a/A) = \text{tp}(b/A)$. It is a bounded A -type-definable equivalence relation.

REMARK 3. $a \xrightarrow{\text{Ls}}_A b \Rightarrow a \xrightarrow{\text{KP}}_A b \Rightarrow a \text{ such that } {}_A b \Rightarrow a \equiv_A b.$

DEFINITION 4. We define the *distance over A*, $d_A(a, b) \in \omega \cup \{\infty\}$, of two tuples a, b of the same length. There are three cases. If $a = b$ we set $d_A(a, b) = 0$. If there is a natural number $n \geq 1$ for which there are infinite A -indiscernible sequences I_1, \dots, I_n and tuples a_1, \dots, a_{n+1} such that $a = a_1, b = a_{n+1}$ and $a_i, a_{i+1} \in I_i$ for all $i = 1, \dots, n$, we define $d_A(a, b)$ as the least such number n . If $a \neq b$ and there is no such n we put $d_A(a, b) = \infty$.

REMARK 4. $a \xrightarrow{\text{Ls}}_A b$ iff $d_A(a, b) < \infty$.

DEFINITION 5. An automorphism $f \in \text{Aut}(\mathfrak{C}/A)$ is *strong over A* if it is a finite product $f = f_1 \circ \dots \circ f_n$ of automorphisms $f_i \in \text{Aut}(\mathfrak{C}/M_i)$ where each M_i is a model containing A . The strong automorphisms over A form a normal subgroup $\text{Autf}(\mathfrak{C}/A)$ of $\text{Aut}(\mathfrak{C}/A)$.

The group of strong automorphisms was introduced by D. Lascar in [16]. This group acts on the tuples of \mathfrak{C} . The orbits of the action are now called Lascar strong types.

REMARK 5. 1. If $d_A(a, b) \leq 1$, then $a \equiv_M b$ for some model $M \supseteq A$.

2. $a \equiv_M b$ for some $M \supseteq A$ implies $d_A(a, b) \leq 2$.

3. $a \xrightarrow{\text{Ls}}_A b$ iff $f(a) = b$ for some $f \in \text{Autf}(\mathfrak{C}/A)$.

Proof. See Proposition 9.12 and Corollary 9.15 of [4]. □

DEFINITION 6. T is *G-compact over A* if $\xrightarrow{\text{Ls}}_A = \xrightarrow{\text{KP}}_A$ for all possible lengths of tuples.

REMARK 6. The following are equivalent:

1. T is G-compact over A

2. $\xrightarrow{\text{Ls}}_A$ is type-definable over A .

3. There is some $n < \omega$ such that, for all tuples a, b : $a \xrightarrow{\text{Ls}}_A b$ iff $d_A(a, b) \leq n$.

Proof. See Remark 10.16 in [4]. The equivalence with item 3 was first proven by L. Newelski in [17]. □

DEFINITION 7. Let E be a 0-type-definable equivalence relation. A *hyperimaginary of sort E* is an equivalence class a_E . Note that $\text{Aut}(\mathfrak{C}/A)$ acts on the hyperimaginaries of sort E by $f(a_E) = f(a)_E$. The hyperimaginary a_E is *A-bounded* if it has a small orbit in this action. The class of all A -bounded hyperimaginaries is $\text{bdd}(A)$.

Note that whenever E is a bounded 0-type-definable equivalence relation, then $a_E \in \text{bdd}(\emptyset)$. Moreover (see Proposition 15.27 in [4]) if $a_E \in \text{bdd}(\emptyset)$, then one can find some bounded 0-type-definable equivalence relation F such that $a_E = a_F$.

DEFINITION 8. Let A be a class of hyperimaginaries. We write $a \equiv_A b$ to mean that there is an automorphism $f \in \text{Aut}(\mathfrak{C})$ fixing all hyperimaginaries of A and such that $f(a) = b$.

REMARK 7. $a \stackrel{\text{KP}}{\equiv}_A b$ iff $a \equiv_{\text{bdd}(A)} b$.

Proof. We sketch the proof. For details see Proposition 15.21 in [4]. Working in $T(A)$ and using Lemma 15.20 from [4], we may assume $A = \emptyset$. From right to left: clear since $a_{\stackrel{\equiv}{\text{KP}}} \in \text{bdd}(\emptyset)$. For the other direction, one should check that $\equiv_{\text{bdd}(\emptyset)}$ is bounded and 0-type-definable. There is a single hyperimaginary $e \in \text{bdd}(\emptyset)$ such that $\equiv_{\text{bdd}(\emptyset)} = \equiv_e$. Let e be of sort E and let c be such that $c_E = e$. Then \equiv_e is type-definable over c by: $x \equiv_e y \Leftrightarrow \exists z(E(z, c) \wedge xc \equiv_{} yz)$. Since it is invariant over \emptyset and type-definable, it is type-definable over \emptyset . The mapping sending each equivalence class a_{\equiv_e} to $\text{tp}(a/c)$ is one-to-one and therefore \equiv_e is bounded. \square

There is a well-known version of this last result for the case *suchthat* $_A$ in terms of imaginaries:

FACT 1. *asuchthat* $_A b$ iff $a \equiv_{\text{acl}^{\text{eq}}(A)} b$.

2. NIP, simple and NTP₂ theories

DEFINITION 9. The formula $\varphi(x, y) \in L$ has the *tree property* (with respect to $k < \omega$) if there is a tree $(a_s : s \in \omega^{<\omega})$ of tuples a_s such that

- $\{\varphi(x, a_{s \cap n}) : n < \omega\}$ is k -inconsistent for all $s \in \omega^{<\omega}$.
- $\{\varphi(x, a_{f \upharpoonright n}) : n < \omega\}$ is consistent for all $f \in \omega^\omega$.

The theory T is *simple* if no formula has the tree property in T .

Simple theories were first defined by S. Shelah in [19], but the most relevant facts were found later by B. Kim and A. Pillay. There are some expository books and our main reference is [4].

FACT 2. Simple theories are G -compact over any set, with distance 2, that is, in a simple theory for any tuples a, b : $a \stackrel{\text{Ls}}{\equiv}_A b$ iff $d_A(a, b) \leq 2$.

Proof. See Proposition 10.12 of [4]. This was first proven by B. Kim [13]. \square

DEFINITION 10. The formula $\varphi(x, y) \in L$ has the *independence property* if there is a sequence of parameters $(a_n : n < \omega)$ such that for every $X \subseteq \omega$ the following is

consistent:

$$\{\varphi(x, a_n) : n \in X\} \cup \{\neg\varphi(x, a_n) : n \notin X\}$$

The theory is NIP (or *dependent*) if no formula has the independence property in T .

The independence property is studied by S. Shelah in [20] and by B. Poizat in [18]. The most recent developments in the model theory of NIP theories are due to S. Shelah, E. Hrushovski and A. Pillay among others. For an exposition we refer to H. Adler [1] and P. Simon [21].

Both in the definition of NIP and of simplicity we may allow the formula $\varphi(x, y)$ to contain some parameters since they can be added to the nodes a_s of the tree or to the tuples a_n .

REMARK 8. The formula $\varphi(x, y) \in L$ has the independence property iff for some indiscernible sequence $(a_n : n < \omega)$ there is some c such that $\models \varphi(c, a_n)$ iff n is even.

Proof. Assume $\varphi(x, y)$ has the independence property. By compactness we may assume that the sequence $(a_n : n < \omega)$ witnessing the independence property is indiscernible. But $\{\varphi(x, a_{2n}) : n < \omega\} \cup \{\neg\varphi(x, a_{2n+1}) : n < \omega\}$ is consistent.

For the other direction, assume there is an indiscernible sequence $(a_n : n < \omega)$ such that $\{\varphi(x, a_{2n}) : n < \omega\} \cup \{\neg\varphi(x, a_{2n+1}) : n < \omega\}$ is consistent. We claim that $\{\varphi(x, a_n) : n \in X\} \cup \{\neg\varphi(x, a_n) : n \in \omega \setminus X\}$ is consistent for all $X \subseteq \omega$. It is enough to check that for any finite disjoint $X, Y \subseteq \omega$, $\Sigma(x) = \{\varphi(x, a_n) : n \in X\} \cup \{\neg\varphi(x, a_n) : n \in Y\}$ is consistent. Let $m_1 < \dots < m_i$ and $k_1 < \dots < k_j$ be respective enumerations of X and Y and choose even numbers $m'_1 < \dots < m'_i$ and odd numbers $k'_1 < \dots < k'_j$ such that $m_1, \dots, m_i, k_1, \dots, k_j$ and $m'_1, \dots, m'_i, k'_1, \dots, k'_j$ have the same order type. By assumption $\{\varphi(x, a_n) : n = m'_1, \dots, m'_i\} \cup \{\neg\varphi(x, a_n) : n = k'_1, \dots, k'_j\}$ is consistent. By indiscernibility $\Sigma(x)$ is consistent. \square

A theory is stable if and only if it is simple and NIP. An important class of unstable NIP theories are the o-minimal theories.

FACT 3. In an o-minimal theory all automorphisms are strong. If two tuples have the same type over A , then they have the same type over some model containing A . Hence, all o-minimal theories are G -compact over any set, with distance 2.

Proof. See Lemma 24 in [22]. \square

DEFINITION 11. The formula $\varphi(x, y) \in L$ has TP_2 , *the tree property of the second kind*, if there is an array of tuples $(a_{ij} : i, j < \omega)$ and some $k < \omega$ such that:

- $\{\varphi(x, a_{ij}) : j < \omega\}$ is k -inconsistent for every $i < \omega$.
- $\{\varphi(x, a_{if(i)}) : i < \omega\}$ is consistent for every $f : \omega \rightarrow \omega$.

The theory T is TP_2 if some formula has the tree property of the second kind in T , otherwise it is NTP_2 .

LEMMA 1. *If T has TP_2 , then some formula $\varphi(x, y)$ has TP_2 in T with respect to $k = 2$.*

Proof. Start with an array $(a_{i,j} : i, j < \omega)$, a natural number k and a formula $\varphi(x, y)$ witnessing TP_2 . We may assume k is minimal, that is, no array and formula witness TP_2 of T with a smaller number. We may also assume that the rows of the array are mutually indiscernible, in the sense that each row is indiscernible over the other rows. For more details on this see, for instance, Lemma 1.2 of [7]. Consider the set $\{\varphi(x, a_{i,0}), \varphi(x, a_{i,1}) : i < \omega\}$. If it is consistent, then (by indiscernibility of the array) the formula $\psi(x; y_0 y_1) = \varphi(x, y_0) \wedge \varphi(x, y_1)$ and the array $(a_{i,2j} a_{i,2j+1} : i < \omega, j < \omega)$ witness TP_2 with a smaller number. If it is inconsistent, we choose $n < \omega$ such that $\{\varphi(x, a_{i,0}), \varphi(x, a_{i,1}) : i < n\}$ is inconsistent. Then the formula $\psi(x; y_0, \dots, y_{n-1}) = \bigwedge_{i < n} \varphi(x, y_i)$ together with the array $(a_{ni,j}, \dots, a_{n(i+1)-1,j} : i < \omega, j < \omega)$ witnesses TP_2 with $k = 2$. \square

REMARK 9. If T is simple or NIP, it is NTP_2 .

Proof. Assume $\varphi(x, y)$ has TP_2 , witnessed by the array $(a_{ij} : i, j < \omega)$ and the number $k < \omega$. If we put $b_\emptyset = a_{00}$ and $b_s = a_{n+1,s(n)}$ for $s \in \omega^{n+1}$, then the tree $(b_s : s \in \omega^{<\omega})$ witnesses that $\varphi(x, y)$ has the tree property with respect to k and, therefore, T is not simple. On the other hand, by Lemma 1 we can assume that $\varphi(x, y)$ has TP_2 with respect to $k = 2$, and then the sequence $(a_{i0} : i < \omega)$ witnesses that $\varphi(x, y)$ has the independence property. \square

NTP_2 theories were defined by S. Shelah in [20] and they are being systematically investigated in the last few years. See A. Chernikov's exposition in [7].

In section 6 we will show that NIP theories are G -compact over extension bases (to be defined in Section 6), with distance 2. This result is due to E. Hrushovski and A. Pillay (see Lemma 2.9 in [10]) and has been recently generalized by I. Ben-Yaacov and A. Chernikov to all NTP_2 theories, but with distance 3 (see Corollary 3.6 of [3]). It is unknown whether it can be improved to distance 2.

3. Dividing, forking and splitting

DEFINITION 12. Let $\varphi(x, y) \in L$.

1. $\varphi(x, a)$ divides over A if there is an A -indiscernible sequence $(a_i : i < \omega)$ such that $\{\varphi(x, a_i) : i < \omega\}$ is inconsistent and $a \equiv_A a_0$.
2. $\varphi(x, a)$ forks over A if it implies a disjunction of formulas each of which divides over A .
3. A partial type divides (forks) over A if it implies a formula that divides (forks) over A .

Dividing implies forking. B. Kim proved in [12] that in simple theories they coincide (see also Proposition 5.17 in [4]). A basic property of dividing is that a partial type over A does not divide over A . This is false for forking in some cases. But forking has the extension property: if a type $p(x)$ over B does not fork over $A \subseteq B$ and $C \supseteq B$ is given, then some complete extension of $p(x)$ over C does not fork over A . This extension property is sometimes false for dividing.

DEFINITION 13. A type $p(x) \in S(B)$ *splits over $A \subseteq B$* if there is a formula $\varphi(x, y) \in L(A)$ and finite tuples $a \equiv_A b$ such that $\varphi(x, a) \in p(x)$ and $\neg\varphi(x, b) \in p(x)$. If $a \xrightarrow{KP} b$ we say that *KP-splits over A* and if $a \xrightarrow{Ls} b$, we say that *$p(x)$ Lascar splits over A* . If $d_A(a, b) \leq 1$ we say that *$p(x)$ strongly splits over A* . The type $p(x)$ is *finitely satisfiable in A* if every finite subset of $p(x)$ is realized in A . In case A is a model this means that $p(x)$ is a *coheir* of $p \upharpoonright A$.

We will discuss all these splitting notions and their relation to dividing and forking. We only consider the case of complete types and the results should not be extrapolated to formulas.

REMARK 10. In general for all $p(x) \in S(B)$ (over $A \subseteq B$)

$$\text{dividing} \Rightarrow \text{forking} \Rightarrow \text{not finitely satisfiable in } A$$

and

$$\text{strongly splitting} \Rightarrow \text{Lascar splitting} \Rightarrow \text{KP-splitting} \Rightarrow \text{splitting} \Rightarrow \text{not fin. sat. in } A$$

Proof. Only the implications *splitting over $A \Rightarrow$ not finitely satisfiable in A* and *forking over $A \Rightarrow$ not finitely satisfiable in A* need some explanation. Assume $p(x) \in S(B)$ is finitely satisfiable in A . If $p(x)$ forks over A , then $p(x) \vdash \psi_1(x, a_1) \vee \dots \vee \psi_n(x, a_n)$ for some formulas $\psi_i(x, y_i) \in L$ and tuples a_i such that $\psi_i(x, a_i)$ divides over A . By finite satisfiability some $\psi_i(x, a_i)$ is satisfied in A , which is incompatible with dividing over A .

If now $p(x)$ splits over A , then for some tuples $a, b \in B$, for some formula $\varphi(x, y) \in L(A)$, $a \equiv_A b$, $\varphi(x, a) \in p(x)$ and $\neg\varphi(x, b) \in p(x)$. Then $\varphi(x, a) \wedge \neg\varphi(x, b)$ is satisfiable in A , contradicting the fact that $a \equiv_A b$. \square

We can obtain better results making B large over A or making A a model.

DEFINITION 14. A model N is ω -saturated over $A \subseteq N$ if for every finite tuple $b \in N$ every n -type over Ab is realized in N .

PROPOSITION 1. *If N is ω -saturated over $A \subseteq N$, then for all $p(x) \in S(N)$ (over A):*

$$\text{dividing} = \text{forking} \Rightarrow \text{strongly splitting} = \text{Lascar splitting} \Rightarrow \text{KP-splitting} \Rightarrow \text{splitting}$$

Proof. $\text{forking} \Rightarrow \text{dividing}$. Assume $p(x) \in S(N)$ and $p(x) \vdash \psi_1(x, a_1) \vee \dots \vee \psi_n(x, a_n)$ where each $\psi_i(x, a_i)$ divides over A . There is some tuple $a \in N$ and some formula $\varphi(x, y) \in L$ such that $\varphi(x, a) \in p(x)$ and $\varphi(x, a) \vdash \psi_1(x, a_1) \vee \dots \vee \psi_n(x, a_n)$. By saturation of N , we can find $b_1, \dots, b_n \in N$ such that $a_1, \dots, a_n \equiv_{Aa} b_1, \dots, b_n$. Then $\varphi(x, a) \vdash \psi_1(x, b_1) \vee \dots \vee \psi_n(x, b_n)$ and each $\psi_i(x, b_i)$ divides over A . It follows that $\psi_i(x, b_i) \in p(x)$ for some i , and this implies that $p(x)$ divides over A .

$\text{dividing} \Rightarrow \text{strongly splitting}$. Assume the formula $\varphi(x, a) \in p(x)$ divides over A , witnessed by the A -indiscernible sequence $(a_i : i < \omega)$. We can assume $a = a_0$. Let $n < \omega$ be such that $\{\varphi(x, a_i) : i \leq n\}$ is inconsistent and find, by saturation, some tuples $a'_1, \dots, a'_n \in N$ such that $a_1, \dots, a_n \equiv_{Aa} a'_1, \dots, a'_n$. Since $p(x)$ is complete and consistent, $\neg\varphi(x, a'_i) \in p(x)$ for some $i \leq n$. Since $d_A(a, a'_i) \leq 1$, this shows that $p(x)$ strongly splits over A .

$\text{Lascar splitting} \Rightarrow \text{strongly splitting}$. Assume $p(x) \in S(N)$ does not strongly split over A , $\varphi(x, y) \in L(A)$, $\varphi(x, a) \in p(x)$, b is a tuple of N and $b \stackrel{\text{Ls}}{\equiv}_A a$. We want to check that $\varphi(x, b) \in p(x)$. For some $n < \omega$, $d_A(a, b) \leq n$. Choose tuples a_1, \dots, a_{n+1} and infinite indiscernible sequences I_1, \dots, I_n such that $a = a_1, b = a_{n+1}$ and $a_i, a_{i+1} \in I_i$ for all i . By saturation of N , there are $b_1, \dots, b_n \in N$ such that $a_1, \dots, a_n \equiv_{Aab} b_1, \dots, b_n$. Then $a = b_1, b = b_n$ and $d_A(b_i, b_{i+1}) \leq 1$ for all i . Inductively we see that $\varphi(x, b_i) \in p(x)$ for all i . Hence $\varphi(x, b) \in p(x)$. \square

PROPOSITION 2. *If $A = M$ is a model and $M \subseteq B$, then for any $p(x) \in S(B)$ (over M):*

$$\text{Lascar splitting} = \text{KP-splitting} = \text{splitting} \Rightarrow \text{not coheir}$$

Proof. We need to check $\text{splitting over } M \Rightarrow \text{Lascar splitting over } M$. But this is clear since $a \equiv_M b$ implies $a \stackrel{\text{Ls}}{\equiv}_M b$ since any automorphism $f \in \text{Aut}(\mathfrak{C}/M)$ is strong. \square

If T is NIP we can obtain more information:

PROPOSITION 3. *Assume T is NIP. Strongly splitting implies dividing and Lascar splitting implies forking. Hence for any $p(x) \in S(B)$ (over $A \subseteq B$):*

$$\begin{array}{ccccccc} \text{strongly splitting} & \Rightarrow & \text{Lascar splitting} & \Rightarrow & \text{KP-splitting} & \Rightarrow & \text{splitting} \\ \Downarrow & & \Downarrow & & & & \Downarrow \\ \text{dividing} & \Rightarrow & \text{forking} & & & & \text{not finitely sat. in } A \end{array}$$

If N is ω -saturated over A then for all $p(x) \in S(N)$ (over A):

$$\text{strongly splitting} = \text{dividing} = \text{forking} = \text{Lascar splitting} = \text{KP-splitting} \Rightarrow \text{splitting}$$

If $A = M \subseteq B$ is a model, then for all $p(x) \in S(B)$ (over M):

$$\begin{array}{ccccccc} \text{strongly splitting} & \Rightarrow & \text{Lascar splitting} & = & \text{KP-splitting} & = & \text{splitting} \\ \Downarrow & & \Downarrow & & & & \Downarrow \\ \text{dividing} & = & \text{forking} & & & & \text{not coheir} \end{array}$$

Hence, in a NIP theory if N is ω -saturated over a model $M \subseteq N$, for any $p(x) \in S(N)$ all these notions coincide, although they are stronger than not being a coheir.


Proof. Let T be NIP. We first prove $\text{strongly splitting over } A \Rightarrow \text{dividing over } A$. Assume $p(x) \in S(B)$ does not divide over $A \subseteq B$ but it strongly splits over A . For some tuples $a, b \in B$, for some formula $\varphi(x, y) \in L(A)$, $d_A(a, b) \leq 1$, $\varphi(x, a) \in p(x)$ and $\neg\varphi(x, b) \in p(x)$. We may assume that there is an A -indiscernible sequence $(a_i : i < \omega)$ with $a_0 = a$ and $a_1 = b$. Then $(a_{2i}a_{2i+1} : i < \omega)$ is A -indiscernible and, since $\varphi(x, a) \wedge \neg\varphi(x, b)$ does not divide over A , $\{\varphi(x, a_{2i}) \wedge \neg\varphi(x, a_{2i+1}) : i < \omega\}$ is consistent. Let c realize this set of formulas. Then $\models \varphi(c, a_i)$ iff i is even. By Remark 8, this implies that $\varphi(x, y)$ has the independence property.

Lascar splitting over $A \Rightarrow$ forking over A . Assume $p(x) \in S(B)$ does not fork over $A \subseteq B$ and choose some model $N \supseteq B$ which is ω -saturated over A . There is an extension $q(x) \in S(N)$ of $p(x)$ that does not fork over A . Hence $q(x)$ does not divide over A and (by NIP) it does not strongly split over A . Hence $q(x)$ does not Lascar split over A and the same can be said of $p(x)$.


The proof of $KP\text{-splitting over } A \Rightarrow \text{Lascar splitting over } A$ for $p(x) \in S(N)$ and N ω -saturated over $A \subseteq N$ is postponed to the last section. See Proposition 5.

The implication $\text{forking over } M \Rightarrow \text{dividing over } M$ holds more generally for all NTP₂ theories. This was first proven by A. Chernikov and I. Kaplan in [8]. For a simplified proof see [2]. \square

The following diagram summarizes the implications in the general case:



If moreover T is NIP, the diagram is as follows:



4. Product of types

DEFINITION 15. Following Lascar [16], we say that a set B is *complete* over $A \subseteq B$ if every n -type over A is realized in B . It is the right assumption to guarantee the existence of nonsplitting extensions.

PROPOSITION 4. *If B is complete over $A \subseteq B$, $p(x) \in S(B)$ does not split over A and $B \subseteq C$, then there is a unique type $q(x) \in S(C)$ extending $p(x)$ that does not split over A .*

Proof. We define:

$$q(x) = p(x) \cup \{\varphi(x, a) : \varphi(x, y) \in L(A), a \in C \text{ and } \varphi(x, a') \in p(x) \text{ for some } a' \equiv_A a\}.$$

The type should extend $q(x)$. Since B is complete over A , $q(x)$ is a complete type over C (if consistent). This proves the uniqueness. We now check the consistency of $q(x)$. Assume, searching for a contradiction, that $p(x) \vdash \neg\varphi_1(x, a_1) \vee \dots \vee \neg\varphi_n(x, a_n)$ where $a_i \in C$, $\varphi_i(x, y_i) \in L(A)$, $a_i \equiv_A a'_i$ and $\varphi_i(x, a'_i) \in p(x)$ for each i . Choose a formula $\theta(x, b) \in p(x)$ which implies the disjunction and choose tuples b', a''_1, \dots, a''_n in B such that $b', a''_1, \dots, a''_n \equiv_A b, a_1, \dots, a_n$. Then $\theta(x, b') \in p(x)$ and $\theta(x, b') \vdash \neg\varphi_1(x, a''_1) \vee \dots \vee \neg\varphi_n(x, a''_n)$. For some i , $\neg\varphi_i(x, a''_i) \in p(x)$, which contradicts non-splitting of $p(x)$ over A since $a''_i \equiv_A a_i$. \square

DEFINITION 16. We will denote by $p|_A C$ the unique nonsplitting extension of $p(x) \in S(B)$ over C as in the previous proposition. See the next remark to evaluate the dependency on A .

REMARK 11. Assume B is AA' -complete and $p(x) \in S(B)$ does not split over A nor over A' . Then for every $C \supseteq B$, $p|_A C = p|_{A'} C$.

Proof. Let $\varphi(x, y) \in L(A)$, $a \in C$ and $\varphi(x, a) \in p|_A C$. Choose a tuple $b \in B$ such that $a \equiv_{AA'} b$. Since $a \equiv_A b$, $\varphi(x, b) \in p \subseteq p|_{A'} C$. Since $a \equiv_{A'} b$, $\varphi(x, a) \in p|_{A'} C$. \square

REMARK 12. If $A \subseteq B \subseteq C \subseteq D$, B is complete over A and $p(x) \in S(B)$ does not split over A , then:

1. $(p|_A C)|_A D = p|_A D$
2. If the sequence $(a_i : i < \omega)$ is such that $a_i \models p|_A B a_{<i}$ for all $i < \omega$, then $(a_i : i < \omega)$ is B -indiscernible.

Proof. 1. Clear, since $(p|_A C)|_A D$ extends p and does not split over A .

2. By induction on $n < \omega$ we prove that for all $i_0 < \dots < i_n < \omega$, $a_{i_0}, \dots, a_{i_n} \equiv_B a_0, \dots, a_n$. The case $n = 0$ is obvious. For the case $n + 1$, assume $\varphi(x_0, \dots, x_{n+1}) \in L(B)$, $i_0 < \dots < i_{n+1}$ and $\models \varphi(a_{i_0}, \dots, a_{i_{n+1}})$. Then $\varphi(a_{i_0}, \dots, a_{i_n}, x) \in p|_A B a_{<i_{n+1}}$ and by nonsplitting and the inductive hypothesis, $\varphi(a_0, \dots, a_n, x) \in p|_A B a_{<n+1}$. Hence $\models \varphi(a_0, \dots, a_{n+1})$. \square

DEFINITION 17. Let $B \supseteq A$ be complete over A and assume $q(y) \in S_y(B)$ does not split over A . For any $p(x) \in S_x(B)$ the *product* $p \otimes_A q$ is the only type in $S_{xy}(B)$ such that for all a, b :

$$ab \models p \otimes_A q \Leftrightarrow a \models p \text{ and } b \models q|_A B a$$

Note that this is independent of the choice of a, b : if a', b' is another choice, then $ab \equiv_B a'b'$. Sometimes the product is defined in the reverse order: one assumes $p(x)$ does not split over A , takes $b \models q$ and $a \models p|_A B b$ and sets $p \otimes_A q = \text{tp}(ab/B)$.

LEMMA 2. Assume B is complete over $A \subseteq B$ and $p(x), q(y), r(z) \in S(B)$ and $q(y), r(z)$ do not split over A . Then:

1. $p \otimes_A q$ and $q \otimes_A r$ do not split over A .
2. $p \otimes_A (q \otimes_A r) = (p \otimes_A q) \otimes_A r$.
3. For any $C \supseteq B$, $(q \otimes_A r)|_A C = q|_A C \otimes_A r|_A C$

Proof. 1. We check this for $p \otimes_A q$. Assume $\varphi(x, y, u) \in L(A)$, c, c' are tuples of B such that $c \equiv_A c'$, $ab \models p \otimes_A q$ and $\models \varphi(a, b, c)$. Then $\varphi(a, y, c) \in q|_A Ba$, a type that does not split over A . Hence $\varphi(a, y, c') \in q|_A Ba$ and $\models \varphi(a, b, c')$.

2. If $a \models p$, $b \models q|_A Ba$ and $c \models r|_A Bab$, then $ab \models p \otimes_A q$ and $abc \models (p \otimes_A q) \otimes_A r$.
3. Clear, since $q|_A C \otimes_A r|_A C$ is an extension of $q \otimes_A r$ that does not split over A . \square

DEFINITION 18. If B is complete over $A \subseteq B$ and $p(x) \in S(B)$ does not split over A , the *power* $p^{(n)_A}$ is defined as the n -times iterated product $p \otimes_A \dots \otimes_A p$. More generally, we can define for any ordinal α the power $p^{(\alpha)_A}$ as the type over B of a sequence $(a_i : i < \alpha)$ such that $a_i \models p|_A Ba_{<i}$ for all $i < \alpha$. These types do not split over A .

Being complete over a subset A is a weaker condition than being a model ω -saturated over A . However, forking still implies splitting in this situation.

REMARK 13. Let B be complete over $A \subseteq B$ and assume $p(x) \in S(B)$ forks over A . Then $p(x)$ splits over A .

Proof. Suppose $p(x)$ does not split over A . Let $N \supseteq B$ be ω -saturated over A . Let $q(x) = p(x)|_A N$. Since $q(x) \in S(N)$ does not split over A , by Proposition 1 it does not fork over A . Then $p(x)$ does not fork over A , a contradiction. \square

A similar treatment of Lascar splitting is possible. We can define B to be *Lascar complete* over $A \subseteq B$ if for every finite tuple a there is some $a' \in B$ such that $a' \stackrel{\text{Ls}}{\equiv}_A a$. An analogous of Proposition 4 holds: if B is Lascar complete over B and $C \supseteq B$, any type $p(x) \in S(B)$ that does not Lascar split over A has a unique extension $q(x) \in S(C)$ that does not Lascar split over A . Using this we can prove, as in Remark 13, that if $p(x) \in S(B)$ forks over $A \subseteq B$ and B is Lascar complete over A , then $p(x)$ Lascar splits over A .

5. Global types

DEFINITION 19. A *global type* is a type $\mathfrak{p}(x)$ over the monster model \mathfrak{C} . Every automorphism $f \in \text{Aut}(\mathfrak{C})$ moves $\mathfrak{p}(x)$ to some *conjugate*

$$\mathfrak{p}^f(x) = \{\varphi(x, f(a)) : \varphi(x, y) \in L, \varphi(x, a) \in \mathfrak{p}(x)\}$$

We say that $\mathfrak{p}(x)$ is *A-invariant* if $\mathfrak{p}^f(x) = \mathfrak{p}(x)$ for all $f \in \text{Aut}(\mathfrak{C}/A)$. Similarly, we say that it is *KP-invariant over A* or *bdd(A)-invariant* if it is fixed under the action of $\text{Aut}(\mathfrak{C}/\text{bdd}(A))$ and we say that it is *Lascar invariant over A* if it is fixed under the action of $\text{Aut}^*(\mathfrak{C}/A)$.

- REMARK 14.
1. $\mathfrak{p}(x)$ is *A-invariant* iff it does not split over *A*.
 2. $\mathfrak{p}(x)$ is *bdd(A)-invariant* iff it does not KP-split over *A*.
 3. $\mathfrak{p}(x)$ is *Lascar invariant over A* iff it does not Lascar split over *A*.

REMARK 15.

1. Since the monster model is ω -saturated over *A*, for any global type $\mathfrak{p}(x)$, over any small set *A*:

$$\text{dividing} = \text{forking} \Rightarrow \text{strongly splitting} = \text{Lascar splitting}$$

2. If *A* is a model, then additionally:

$$\text{invariant} = \text{KP-invariant} = \text{Lascar invariant}$$

3. In a NIP theory, $\mathfrak{p}(x)$ does not divide over *A* iff it does not fork over *A* iff it is Lascar invariant over *A* iff it is KP-invariant over *A*. If additionally *A* is a model, these conditions are also equivalent to being *A-invariant*.

Recall that a global type $\mathfrak{p}(x)$ is *A-definable* if for each $\varphi(x, y) \in L$ the set $\{a : \varphi(x, a) \in \mathfrak{p}(x)\}$ is definable over *A*. The type is definable if it is *A-definable* for some small set *A*.

REMARK 16. Assume $\mathfrak{p}(x)$ is definable. Then $\mathfrak{p}(x)$ is definable over *A* iff it is *A-invariant*.

Proof. Let $X_\varphi = \{a : \varphi(x, a) \in \mathfrak{p}(x)\}$. If \mathfrak{p} is *A-definable*, then $f(X_\varphi) = X_\varphi$ and hence $\mathfrak{p}^f = \mathfrak{p}$ for every $f \in \text{Aut}(\mathfrak{C}/A)$. For the other direction use Remark 1. \square

DEFINITION 20. A global type $\mathfrak{p}(x)$ is called *invariant* if it is *A-invariant* for some set *A*.

REMARK 17. If $\mathfrak{p}(x)$ is *A-invariant*, $B \supseteq A$ is complete over *A*, and $p(x) = \mathfrak{p} \upharpoonright B$, then $p(x)|_A \mathfrak{C} = \mathfrak{p}(x)$.

Proof. $\mathfrak{p}(x)$ is an extension of $p(x)$ that does not split over *A*. \square

Let \mathfrak{q} be an invariant global type. There are different ways to define the product $\mathfrak{p} \otimes \mathfrak{q}$. One option is to step outside the monster model \mathfrak{C} and work in another monster model \mathfrak{C}' extending \mathfrak{C} where every type over \mathfrak{C} and any small subset of \mathfrak{C}' is realized. Then we can realize $a \models \mathfrak{p}$ and $b \models \mathfrak{q}|_{\mathfrak{C}a}$ in \mathfrak{C}' and define $\mathfrak{p} \otimes \mathfrak{q} = \text{tp}(ab/\mathfrak{C})$. By Remark 11, this is independent of the choice of the small set *A* over which \mathfrak{q} is invariant.

Another equivalent possibility is to choose a set $B \supseteq A$ which is complete over A and define $\mathfrak{p} \otimes \mathfrak{q} = (\mathfrak{p} \upharpoonright B \otimes_A \mathfrak{q} \upharpoonright B)|_A \mathfrak{C}$. There is a third option, described in the next result.

REMARK 18. Let $\mathfrak{q}(x)$ and $\mathfrak{q}(y)$ be global types, let A be a small set and assume $\mathfrak{q}(y)$ is A -invariant. For each $C \supseteq A$ let $r_C(x, y) = \text{tp}(ab/C)$, where $a \models \mathfrak{p} \upharpoonright C$ and $b \models \mathfrak{q} \upharpoonright Ca$. Note that r_C is well-defined independently of the choice of a and b . Since $C \subseteq C'$ implies $r_C \subseteq r_{C'}$ the type $\mathfrak{r}(x, y) = \bigcup_{C \supseteq A} r_C$ is a global type. Then $\mathfrak{r} = \mathfrak{p} \otimes \mathfrak{q}$.

Proof. We first check that $r_C(x, y)$ is well-defined. Assume $a_i \models \mathfrak{p} \upharpoonright C$ and $b_i \models \mathfrak{q} \upharpoonright Ca_i$ for $i = 1, 2$. Choose $f \in \text{Aut}(\mathfrak{C}/C)$ such that $f(a_1) = a_2$ and let $b = f(b_1)$. Since $\mathfrak{q}^f = \mathfrak{q}$, $b \models \mathfrak{q} \upharpoonright Ca_2$ and hence $a_1 b_1 \equiv_C a_2 b \equiv_C a_2 b_2$.

For the rest it is enough to show that $r_B = \mathfrak{p} \upharpoonright B \otimes_A \mathfrak{q} \upharpoonright B$ for every $B \supseteq A$ complete over A . And this is clear, because $(\mathfrak{q} \upharpoonright B)|_A Ba = \mathfrak{q} \upharpoonright Ba$ for any $a \models \mathfrak{p} \upharpoonright B$. \square

More generally, one can define in a similar way the power $\mathfrak{p}^{(\alpha)} = \mathfrak{p}^{(\alpha)}(x_i : i < \alpha)$ for any invariant type \mathfrak{p} and any ordinal α . We set $\mathfrak{p}^{(\alpha+1)}(x_i : i \leq \alpha) = \mathfrak{p}^{(\alpha)}(x_i : i < \alpha) \otimes \mathfrak{p}(x_\alpha)$ and for limit α , $\mathfrak{p}^{(\alpha)}(x_i : i < \alpha) = \bigcup_{\beta < \alpha} \mathfrak{p}^{(\beta)}(x_i : i < \beta)$.

REMARK 19. 1. If \mathfrak{q} is A -invariant, then $\mathfrak{p} \otimes \mathfrak{q}$ is A -invariant.

2. If $\mathfrak{q}, \mathfrak{r}$ are invariant types, then $\mathfrak{p} \otimes (\mathfrak{q} \otimes \mathfrak{r}) = (\mathfrak{p} \otimes \mathfrak{q}) \otimes \mathfrak{r}$.

3. If \mathfrak{p} is A -invariant, then $\mathfrak{p}^{(\alpha)}$ is A -invariant for every ordinal α .

4. If \mathfrak{p} is A -invariant, then any realization $(a_i : i < \alpha)$ of the power $\mathfrak{p}^{(\alpha)}$ is an A -indiscernible sequence.

Proof. By Lemma 2. \square

DEFINITION 21. A *Morley sequence* in \mathfrak{p} over A is a sequence $(a_i : i < \alpha)$ such that $a_i \models \mathfrak{p} \upharpoonright Aa_{<i}$ for all $i < \alpha$.

REMARK 20. Let \mathfrak{p} be A -invariant. A sequence $(a_i : i < \alpha)$ is a Morley sequence in \mathfrak{p} over A iff it realizes $\mathfrak{p}^{(\alpha)} \upharpoonright A$.

Proof. Induction on α . \square

LEMMA 3. $\mathfrak{p}(x)$ is Lascar invariant over A if and only if it is M -invariant for every model $M \supseteq A$.

Proof. If $A \subseteq M$, then $\text{Aut}(\mathfrak{C}/M) \subseteq \text{Aut}(\mathfrak{C}/A)$. On the other hand, every strong automorphism $f \in \text{Aut}(\mathfrak{C}/A)$ is of the form $f = f_1 \circ \dots \circ f_n$ where for every i , $f_i \in \text{Aut}(\mathfrak{C}/M_i)$ for some model $M_i \supseteq A$. \square

In particular, Lascar invariant types are invariant. Hence product and powers are well-defined.

REMARK 21. If \mathfrak{p} is Lascar invariant over A , then $\mathfrak{p}^{(\alpha)}$ is Lascar invariant over A for every ordinal α .

Proof. By Lemma 3 and Remark 19. \square

6. *G*-compactness in NIP theories

LEMMA 4. *Assume the global type $\mathfrak{p}(x)$ is Lascar invariant over A . Any realization $(a_i : i < \omega)$ of $\mathfrak{p}^{(\omega)} \upharpoonright A$ is A -indiscernible.*

Proof. Assume $(a_i : i < \omega) \models \mathfrak{p}^{(\omega)} \upharpoonright A$ and choose a model $M \supseteq A$. Then \mathfrak{p} is M -invariant. Let $(a'_i : i < \omega) \models \mathfrak{p}^{(\omega)} \upharpoonright M$. By Remark 19 $(a'_i : i < \omega)$ is M -indiscernible and hence A -indiscernible. Since $(a_i : i < \omega) \equiv_A (a'_i : i < \omega)$, clearly $(a_i : i < \omega)$ is A -indiscernible. \square

THEOREM 4 (Hrushovski-Pillay). *Let $\mathfrak{p}(x)$ be a global type, Lascar invariant over A . If $a \stackrel{\text{Ls}}{\equiv}_A b$ are realizations of $\mathfrak{p} \upharpoonright A$, then there is some realization $(a_i : i < \omega)$ of $\mathfrak{p}^{(\omega)} \upharpoonright A$ such that the sequences a, a_0, a_1, \dots and b, b_0, b_1, \dots are both A -indiscernible. Hence $d_A(a, b) \leq 2$.*

Proof. We can assume $a \models \mathfrak{p} \upharpoonright M$ for some model $M \supseteq A$. The reason is that we can choose a model $N \supseteq A$, some $c \models \mathfrak{p} \upharpoonright N$ and some automorphism $f \in \text{Aut}(\mathfrak{C}/A)$ sending c to a . Then put $M = f(N)$, and let $\mathfrak{q} = \mathfrak{p}^f$ be the A -conjugate of \mathfrak{p} by f . Notice that $a \models \mathfrak{q} \upharpoonright M$ and $\mathfrak{p}^{(\omega)} \upharpoonright A = \mathfrak{q}^{(\omega)} \upharpoonright A$.

Now let $(a_i : i < \omega)$ be a realization of $\mathfrak{p}^{(\omega)} \upharpoonright Mab$. It follows that the sequence a, a_0, a_1, \dots is a Morley sequence in \mathfrak{p} over M and therefore it is M -indiscernible and A -indiscernible. We claim that for each $n < \omega$, $a \equiv_{Aa_0, \dots, a_n} b$, which implies that also b, b_0, b_1, \dots is A -indiscernible. Let $\varphi(x, y_0, \dots, y_n) \in L(A)$. Since $\mathfrak{p}^{(\omega)}$ does not Lascar split over A ,

$$\models \varphi(a, a_0, \dots, a_n) \Leftrightarrow \varphi(a, y_0, \dots, y_n) \in \mathfrak{p}^{(\omega)} \Leftrightarrow \varphi(b, y_0, \dots, y_n) \in \mathfrak{p}^{(\omega)} \Leftrightarrow \models \varphi(b, a_0, \dots, a_n).$$

\square

COROLLARY 1. *If T is NIP and $p(x) \in S(A)$ does not fork over A , then for any realizations a, b of $p(x)$: $a \stackrel{\text{Ls}}{\equiv}_A b$ if and only if $d_A(a, b) \leq 2$.*

Proof. If $p(x) \in S(A)$ does not fork over A , it has a global nonforking extension $\mathfrak{p}(x)$. Since T is NIP, $\mathfrak{p}(x)$ is Lascar invariant over A . \square

An *extension base* is a set A such that no type over A forks over A . It follows that any NIP theory is *G*-compact over extension bases.

LEMMA 5. *Let T be NIP and let $\mathfrak{p}_1(x), \mathfrak{p}_2(x)$ be global types, Lascar invariant over A . If for some model $M \supseteq A$ there is a realization $I = (a_i : i < \omega)$ of $\mathfrak{p}_1^{(\omega)} \upharpoonright M$, such that $\mathfrak{p}_1 \upharpoonright AI = \mathfrak{p}_2 \upharpoonright AI$, then $\mathfrak{p}_1 = \mathfrak{p}_2$.*

Proof. We claim that if $I' = (a_i : i < \alpha)$ is an A -indiscernible sequence extending I , then $I'c$ is also A -indiscernible for any $c \models \mathfrak{p}_1 \upharpoonright AI'$ or $c \models \mathfrak{p}_2 \upharpoonright AI'$. Consider the case $c \models \mathfrak{p}_1 \upharpoonright AI'$. Assume $i_0 < \dots < i_n < \alpha$, $\psi(x_0, \dots, x_n, y) \in L(A)$ and $\models \psi(a_{i_0}, \dots, a_{i_n}, c)$. Then $\psi(a_{i_0}, \dots, a_{i_n}, y) \in \mathfrak{p}_1$. Since \mathfrak{p}_1 does not Lascar-split over A and $a_{i_0} \dots a_{i_n} \stackrel{\text{Ls}}{\equiv}_A a_0 \dots a_n$, $\psi(a_0, \dots, a_n, y) \in \mathfrak{p}_1$. Since $a_{n+1} \models \mathfrak{p}_1 \upharpoonright Ma_0 \dots a_n$, $\models \psi(a_0, \dots, a_n, a_{n+1})$. The case $c \models \mathfrak{p}_2 \upharpoonright AI'$ is similar but uses the assumption $\mathfrak{p}_1 \upharpoonright AI = \mathfrak{p}_2 \upharpoonright AI$.

Now assume $\varphi(x, y) \in L$, $\varphi(x, b) \in \mathfrak{p}_1$ and $\neg\varphi(x, b) \in \mathfrak{p}_2$. Construct $(c_i : i < \omega)$ in such a way that $c_{2i} \models \mathfrak{p}_1 \upharpoonright AIBC_{<2i}$ and $c_{2i+1} \models \mathfrak{p}_2 \upharpoonright AIBC_{<2i+1}$. Note that I is A -indiscernible. By the claim $I^\frown (c_i : i < \omega)$ is also A -indiscernible. Since $\models \varphi(a_{2i}, b)$ and $\models \varphi(a_{2i+1}, b)$, $\varphi(x, y)$ has the independence property, a contradiction. \square

LEMMA 6. *Let T be NIP, let $f \in \text{Aut}(\mathfrak{C}/A)$ and let \mathfrak{p} be a global type which is Lascar invariant over A . Assume that for each $n < \omega$, for each $a \models \mathfrak{p}^{(n)} \upharpoonright A$, $a \stackrel{\text{Ls}}{\equiv}_A f(a)$. Then $\mathfrak{p}^f = \mathfrak{p}$.*

Proof. Choose a model $M \supseteq A$ and let $I = (a_i : i < \omega) \models \mathfrak{p}^{(\omega)} \upharpoonright M$. By Lemma 5 it will suffice to prove $\mathfrak{p} \upharpoonright AI = \mathfrak{p}^f \upharpoonright AI$. Let $\varphi(x, y_0, \dots, y_n) \in L(A)$ and assume $\varphi(x, a_0, \dots, a_n) \in \mathfrak{p}$. Since a_0, \dots, a_n realizes $\mathfrak{p}^{(n+1)} \upharpoonright A$, by assumption $a_0, \dots, a_n \stackrel{\text{Ls}}{\equiv}_A f(a_0), \dots, f(a_n)$. Since \mathfrak{p} does not Lascar split over A , $\varphi(x, f(a_0), \dots, f(a_n)) \in \mathfrak{p}$. This shows that $\mathfrak{p} \upharpoonright AI \supseteq \mathfrak{p}^f \upharpoonright AI$ and the equality. \square

PROPOSITION 5. *Assume T has NIP. Let M be ω -saturated over $A \subseteq M$ and $p(x) \in S(M)$. Then p Lascar splits over A if and only if p KP-splits over A .*

Proof. We only need to prove the direction from right to left. If $p(x)$ does not Lascar split over A , then it does not fork over A and therefore it has a global extension \mathfrak{p} which does not fork over A . Hence \mathfrak{p} is Lascar invariant over A . It is enough to check that \mathfrak{p} is $\text{bdd}(A)$ -invariant. For this purpose, let $f \in \text{Aut}(\mathfrak{C}/\text{bdd}(A))$ and let us prove that $\mathfrak{p}^f = \mathfrak{p}$ using Lemma 6. Let $a \models \mathfrak{p}^{(n)} \upharpoonright A$. Note that $f(a)$ also realizes $\mathfrak{p}^{(n)} \upharpoonright A$ and $a \stackrel{\text{KP}}{\equiv}_A f(a)$. Since $\mathfrak{p}^{(n)}$ does not fork over A , by Corollary 1, $a \stackrel{\text{Ls}}{\equiv}_A f(a)$. \square

REMARK 22. Using the comments made on Lascar completeness after Remark 13 we can easily obtain a sharper version of Proposition 5: If T is NIP and B is Lascar complete over $A \subseteq B$, then for any $p(x) \in S(B)$, p Lascar splits over A if and only if p KP-splits over A .

There are non G -compact theories. The first example was due to M. Ziegler and it is presented in [5]. There are also ω -categorical examples (see [11] and [6]). L. Newelski proved in [17] that in any small theory, for finite tuples $\stackrel{\text{Ls}}{\equiv}_A$ and $\stackrel{\text{KP}}{\equiv}_A$ coincide for any finite set A . In [9] A. Conversano and A. Pillay exhibit a natural example of a non G -compact theory. It is a principal homogeneous space on G , a saturated elementary extension of the universal cover of the group $\text{SL}(2, \mathbb{R})$. This is a NIP example, so in Corollary 1 the assumption that $p(x) \in S(A)$ does not fork over A is essential.

References

- [1] ADLER H., *Introduction to theories without the independence property*. Preprint available at <http://www.logic.univie.ac.at/~adler/docs/nip.pdf>, January 2007.
- [2] ADLER H. *Kim's lemma for NTP₂ theories: a simpler proof of a result by Chernikov and Kaplan*. Rend. Sem. Mat. Univ. Politec. Torino, **72**, 3/4, 121–126.
- [3] BEN-YACOV I. AND CHERNIKOV A. *An independence theorem for NTP₂ theories*. The Journal of Symbolic Logic **79** (2014), 135–153.
- [4] CASANOVAS E. *Simple theories and hyperimaginaries*, vol. 39 of Lecture Notes in Logic, Cambridge University Press, 2011.
- [5] CASANOVAS E., LASCAR D., PILLAY A., AND ZIEGLER M. *Galois groups of first order theories*. Journal of Mathematical Logic **1** (2001), 305–319.
- [6] CASANOVAS E., PELÁEZ R., AND ZIEGLER M. *On many-sorted ω -categorical theories*. Fundamenta Mathematicae **214** (2011), 285–294.
- [7] CHERNIKOV A. *Theories without the tree property of the second kind*. Annals of Pure and Applied Logic **165** (2014), 695–723.
- [8] CHERNIKOV A. AND KAPLAN I. *Forking and dividing in NTP₂ theories*. The Journal of Symbolic Logic **77** (2012) 1–20.
- [9] CONVERSANO A. AND PILLAY A. *Connected components of definable groups and o-minimality I*. Advances in Mathematics **231** (2012), 605–623.
- [10] HRUSHOVSKI E. AND PILLAY A. *On NIP and invariant measures*. Journal of the European Mathematical Society **13** (2011), 1005–1061.
- [11] IVANOV A.A. *A countably categorical theory which is not G-compact*. Siberian Advances in Mathematics **20** (2010), 75–82. Based on arXiv:math/602126.
- [12] KIM B. *Forking in simple unstable theories*. Journal of the London Mathematical Society **57** (1998), 257–267.
- [13] KIM B. *A note on Lascar strong types in simple theories*. The Journal of Symbolic Logic **63** (1998), 926–936.
- [14] KIM B. AND PILLAY A. *Simple theories*. Annals of Pure and Applied Logic **88** (1997), 149–164.
- [15] LASCAR D. *On the category of models of a complete theory*. The Journal of Symbolic Logic **47** (1982), 249–266.
- [16] LASCAR D. *Stability in Model Theory*. Longman Scientific & Technical, Harlow, U.K., 1987.
- [17] NEWELSKI L. *The diameter of a Lascar strong type*. Fundamenta Mathematicae **176** (2):157–170, 2003.
- [18] POIZAT B. *Théories instables*. The Journal of Symbolic Logic **46** (1981), 531–562.
- [19] SHELAH S. *Simple unstable theories*. Annals of Mathematical Logic **19** (1980), 177–203.
- [20] SHELAH S. *Classification Theory*. North Holland P.C., Amsterdam, second edition, 1990.
- [21] SIMON P. *A guide to NIP theories*. To appear in Lecture Notes in Logic, Cambridge University Press. arXiv:1208.3944 [math.LO], August 2012.

- [22] ZIEGLER, M. *Introduction to the Lascar group*. In TENT K., ed. , *Tits Buildings and the Model Theory of Groups*, number 291 in London Mathematical Society Lecture Notes Series, pages 279–298. Cambridge University Press, Cambridge, 2002.

AMS Subject Classification: 03C45

Enrique CASANOVAS,
Departamento de Lógica, Universidad de Barcelona
Montalegre 6, 08001 Barcelona, ESPAÑA
e-mail: e.casanovas@ub.edu

Lavoro pervenuto in redazione il 05.03.2014, e, in forma definitiva, il 05.12.2014