

Alberto Dolcetti and Donato Pertici

DIFFERENTIAL PROPERTIES OF SPACES OF SYMMETRIC REAL MATRICES

Abstract. We study the differential geometric properties of the manifold of non-singular symmetric real matrices endowed with the trace metric; in case of positive definite matrices we describe the full group of isometries.

Keywords. Non-singular symmetric matrices, positive definite matrices, trace metric, symmetric (Semi-)Riemannian spaces, isometries, representations of Lie groups, inner and outer automorphisms.

Introduction.

In this paper we carry on with the study of the Semi-Riemannian submanifolds of the Semi-Riemannian manifold of non-singular real matrices, GL_n , of order $n \geq 2$, endowed with the so-called trace metric g . We started this investigation in [6] (relatively to whole GL_n and SL_n) and in [7] (relatively to O_n and SO_n). In this context, the totally geodesic submanifolds of (GL_n, g) (as SL_n and O_n) seem to have a particular role. For this reason we focus our attention on the manifolds $(GLSym_n(p), g)$ of the non-singular symmetric real matrices of signature $(p, n - p)$ together with their Semi-Riemannian submanifolds $(SLSym_n(p), g)$ of matrices with determinant $(-1)^{n-p}$; indeed also all these manifolds are totally geodesic in (GL_n, g) and so their properties can be deduced from the analogous properties of (GL_n, g) . Furthermore, as particular cases among them, we find (\mathcal{P}_n, g) , the Riemannian manifold of real symmetric positive definite matrices, and its submanifold $(SL\mathcal{P}_n, g)$, where $SL\mathcal{P}_n = \mathcal{P}_n \cap SL_n$. These last manifolds have a remarkable interest in many frameworks, for instance in theory of metric spaces of non-positive curvature and in matrix information geometry, and they have recently been object of many researches; for more information we refer for instance to [24], [25], [5], [16, Ch. XII], [17], [4], [3], [19], [21].

This paper develops the differential-geometric properties of $(GLSym_n(p), g)$, independently of the signature, and therefore includes (\mathcal{P}_n, g) in a more general setting. In fact in Section 2, we consider the manifolds $(GLSym_n(p), g)$ together with their submanifolds $(SLSym_n(p), g)$, as totally geodesic Semi-Riemannian submanifolds of (GL_n, g) . In particular we obtain suitable expressions for geodesics, parallel transport, Riemann tensor, Ricci and scalar curvature and we prove that $(GLSym_n(p), g)$ is isometric to the Semi-Riemannian product $(SLSym_n(p) \times \mathbb{R}, g \times h)$, where h is the euclidean metric.

In Section 3 we resume many properties of the Riemannian manifold (\mathcal{P}_n, g) and of its submanifold $(SL\mathcal{P}_n, g)$. These properties are mostly already known in literature, but we deduce them again as easy consequences of the results of Section 2.

The last Section 4 is devoted to determine and describe geometrically the full group of

the isometries of $(SL\mathcal{P}_n, g)$ (Theorem 4.1) and of (\mathcal{P}_n, g) (Theorem 4.2) (to our knowledge, not yet known in a complete and explicit way). At first we determine the isometries of $(SL\mathcal{P}_n, g)$ by using many arguments from the theory of symmetric Riemannian spaces and from the theory of Lie group representations; then the isometries of (\mathcal{P}_n, g) are deduced from the de Rham decomposition of (\mathcal{P}_n, g) as $(SL\mathcal{P}_n \times \mathbb{R}, g \times h)$. These arguments seem to work only for (\mathcal{P}_n, g) and do not generalize directly to the study of the full group of the isometries of $(GLSym_n(p), g)$. Finally, we interpret many isometries as suitable symmetries inside (\mathcal{P}_n, g) .

After finishing this work, we have found a paper of Lajos Molnár, where he describes the isometries of the manifold of positive definite hermitian matrices H_n endowed with a class of metrics, which includes g (see [20, Thm. 3]). Comparing this with our results, it follows that every isometry of (\mathcal{P}_n, g) is the restriction of an isometry of (H_n, g) . Anyway our methods are different and should clarify the geometric descriptions and make explicit the links with the theory of symmetric Riemannian spaces and with the theory of Lie group representations.

Acknowledgement. We thank Fabio Podestà for many discussions about the matter of this paper and especially for his fundamental help in clarifying to us the tools of the theory of Lie group representations, used here.

We thank also the referee for the suggestions in order to precise some results and to improve the setting of our work in literature.

1. Preliminary facts and recalls

1.1 Notations. In this paper, for every integer $n \geq 2$, we denote

- $M_n = M_n(\mathbb{R})$: the vector space of real square matrices of order n ;
- $Sym_n = Sym_n(\mathbb{R})$ (resp. Sym_n^0): the vector subspace of M_n of symmetric matrices (resp. with trace equal to 0);
- $GL_n = GL_n(\mathbb{R})$ (respectively $GL_n^+ = GL_n^+(\mathbb{R})$ and $SL_n = SL_n(\mathbb{R})$): the multiplicative group of non-degenerate matrices in M_n (respectively with positive determinant and with determinant equal to 1);
- $GLSym_n = GL_n \cap Sym_n = \{A \in GL_n / A = A^T\}$ (A^T is the transpose of A);
- \mathcal{P}_n (respectively $SL\mathcal{P}_n$): the subset of $GLSym_n$ of positive definite matrices (respectively with determinant 1);
- $GLSym_n(p)$ ($0 \leq p \leq n$): the set of matrices of $GLSym_n$ with signature $(p, n-p)$; in particular $GLSym_n(n) = \mathcal{P}_n$;
- $SLSym_n(p)$ ($0 \leq p \leq n$): the set of matrices of $GLSym_n(p)$ with determinant $(-1)^{n-p}$; in particular $SLSym_n(n) = SL\mathcal{P}_n$;
- J_p ($0 \leq p \leq n$): the diagonal matrix $diag(I_p, -I_{n-p})$, where I_h is the identity matrix of order h , with the agreement that $J_n = I_n$ and $J_0 = -I_n$;
- O_n (resp. SO_n): the group of (resp. special) orthogonal matrices;
- $O_n(p)$ (resp. $SO_n(p)$), $0 \leq p \leq n$: the subgroup of matrices $A \in GL_n$ (resp. $A \in GL_n^+$) such that $AJ_pA^T = J_p$ (in particular, if $p = n$, then $O_n(n) = O_n$ and $SO_n(n) = SO_n$);

- for every $n \geq 3$, $\pi_n : Spin_n \rightarrow SO_n$: the *universal covering* of SO_n , where $Spin_n$ is the usual *Spin group*.

For every *connected Lie group* \mathbf{G} , we denote by $Aut(\mathbf{G})$, by $Inn(\mathbf{G})$ and by $Out(\mathbf{G}) = Aut(\mathbf{G})/Inn(\mathbf{G})$, respectively, the group of *automorphisms*, the group of *inner automorphisms* and the quotient group of *outer automorphisms* of \mathbf{G} .

Since $Spin_n$ is compact, simply connected, simple Lie group, $Out(Spin_n)$ is isomorphic to the group of symmetries of the *Dynkin diagram* of its *Lie algebra* (see for instance [28, Thm. 8.11.3] and [23, pag. 49]). If $n = 2m + 1 \geq 3$ the corresponding Dynkin diagram is B_m and if $n = 2m \geq 4$ it is D_m (see for instance [28, Thm. 8.9.12]); as a consequence $Out(Spin_{2m+1})$ is trivial for every m , while $Out(Spin_{2m}) \simeq \mathbb{Z}_2$ for $m \neq 4$ and $Out(Spin_8) \simeq Dih_3$ (the *dihedral group*).

As usual, the *commutator* of $A, B \in M_n$ is $[A, B] = AB - BA$.

For every $A \in GL_n$, A^{-T} denotes the matrix $(A^T)^{-1} = (A^{-1})^T$.

For every $1 \leq i, j \leq n$, $E^{(i,j)}$ denotes the matrix in M_n whose (h, k) -entry is 1 if $(h, k) = (i, j)$ and 0 otherwise.

For every $A \in M_n$ we denote the *exponential mapping* by $e^A = exp(A) = I_n + \sum_{i=1}^{+\infty} \frac{A^i}{i!}$.

We define a C^∞ -tensor g of type $(0, 2)$ on GL_n , by

$$g_A(V, W) = tr(A^{-1}VA^{-1}W)$$

(tr indicates the *trace* of a matrix). We call *trace metric* the metric induced by g and will denote by g also its restriction to every submanifold of GL_n .

For every $A \in M_n$ and every $C \in GL_n$ we denote by Γ_C the mapping:

- $\Gamma_C(A) = CAC^T$ (*congruence by C*);

if, moreover, $A \in GL_n$, we denote by φ , by φ_C and by ψ the following mappings:

- $\varphi(A) = A^{-1}$ (*inversion*);

- $\varphi_C(A) = (\Gamma_C \circ \varphi)(A) = CA^{-1}C^T$;

- $\psi(A) = |det(A)|^{-2/n}A$.

In particular, for every $A \in GL_n$, we have: $det(\psi(A)) = \frac{1}{det(A)}$,

$(\varphi \circ \psi)(A) = (\psi \circ \varphi)(A) = |det(A)|^{2/n}A^{-1}$; finally φ , ψ , $\varphi \circ \psi$ have always period 2 and, if $C \in GLSym_n$, then also φ_C has period 2 and $\varphi_C(C) = C$.

1.2 Remarks. We recall some facts which are known or easy to check.

a) The sets $GLSym_n(p)$ are the $(n + 1)$ (open) connected components of $GLSym_n$.

b) The mapping: $(C, A) \mapsto \Gamma_C(A) = CAC^T$, is a left action of the group GL_n on every $GLSym_n(p)$.

The action of GL_n on \mathcal{P}_n has been already described in [5, p.314 ff.].

The quotient group $GL_n/\{\pm I_n\}$ acts *effectively* by congruence on every $GLSym_n(p)$. Indeed, arguing on the matrices $E^{(i,j)} + E^{(j,i)}$, on Sym_n it is simple to check that $\Gamma_C = \Gamma_{C'}$ if and only if $C = \pm C'$ and this suffices to conclude, being Sym_n the tangent space to $GLSym_n(p)$ at any point.

If $A \in GLSym_n(p)$, there exists a matrix $C \in GL_n$ such that $\Gamma_C(A) = CAC^T = J_p$. This implies that, for every p , both GL_n and GL_n^+ act transitively on $GLSym_n(p)$.

Moreover $O_n(p)$ and $SO_n(p)$ are the *isotropy* subgroups at J_p with respect to these actions. Hence $GLSym_n(p)$ is diffeomorphic to both *homogeneous manifolds* $GL_n/O_n(p)$ and $GL_n^+/SO_n(p)$. In particular $\mathcal{P}_n \simeq GL_n/O_n \simeq GL_n^+/SO_n$ (see for instance [27, Thm. 3.62]).

c) Since SL_n acts transitively by congruence on $SLSym_n(p)$, as above we get that $SLSym_n(p)$ is a submanifold of $GLSym_n(p)$, diffeomorphic to the homogeneous manifold $SL_n/SO_n(p)$. In particular $SL\mathcal{P}_n \simeq SL_n/SO_n$.

d) An *isometry* between two Semi-Riemannian manifolds is a diffeomorphism between them preserving metric tensors. If $(\widehat{\mathcal{M}}, \widehat{g})$ is any Semi-Riemannian manifold we denote by $I(\widehat{\mathcal{M}}, \widehat{g})$ the set of isometries of $(\widehat{\mathcal{M}}, \widehat{g})$.

It is known that $I(\widehat{\mathcal{M}}, \widehat{g})$ has the structure of Lie group (see for instance [22, Ch. 9 Thm. 32]) and we will denote by $I^0(\widehat{\mathcal{M}}, \widehat{g})$ its connected component containing the identity.

We end this section by recalling the following

1.3 Proposition. *a) (GL_n, g) is a homogeneous geodesically complete Semi-Riemannian manifold of signature $(\frac{n(n+1)}{2}, \frac{n(n-1)}{2})$, whose geodesics are the curves:*

$t \mapsto Ke^{tC}$ for every $C \in M_n$ and $K \in GL_n$.

b) The Levi-Civita connection ∇ of (GL_n, g) is

$$(\nabla_X \mathcal{Y})_K = (X(\mathcal{Y}))_K - \frac{1}{2}(\mathcal{X}_K K^{-1} \mathcal{Y}_K + \mathcal{Y}_K K^{-1} \mathcal{X}_K),$$

for every $K \in GL_n$ and for all tangent vector fields X, \mathcal{Y} of class C^∞ on GL_n ,

and the Riemann curvature tensor of type $(0, 4)$ of (GL_n, g) is

$$R_{XYZW}(K) = \frac{1}{4}tr([K^{-1}X, K^{-1}Y][K^{-1}Z, K^{-1}W]),$$

for every $K \in GL_n$ and every $X, Y, Z, W \in M_n = T_K(GL_n)$.

c) (GL_n^+, g) is a symmetric Semi-Riemannian manifold and among its isometries there are the congruences and the inversion φ .

Proof. See [6, Prop. 1.1, Thm. 2.1, Prop. 3.1 and Prop. 1.2]. □

2. The Semi-Riemannian manifolds $(GLSym_n(p), g)$

2.1 Proposition. *a) $(GLSym_n(p), g)$ is a homogeneous Semi-Riemannian submanifold of (GL_n, g) with signature $(\frac{n(n+1)}{2} - p(n-p), p(n-p))$ for every $p = 0, \dots, n$.*

An orthonormal basis with respect to g_{J_p} of the tangent space $T_{J_p}(GLSym_n(p)) = Sym_n$ is

$$\mathcal{B} = \{E^{(i,i)} : i = 1, \dots, n\} \cup \{S^{(i,j)} = \frac{E^{(i,j)} + E^{(j,i)}}{\sqrt{2}} : 1 \leq i < j \leq n\}.$$

The time-like vectors are the vectors $S^{(i,j)}$ with $1 \leq i \leq p, p+1 \leq j \leq n$, the remaining vectors are space-like.

b) For every $p = 0, \dots, n$, the mapping $A \mapsto -A$ is an isometry between the Semi-Riemannian manifolds $(GLSym_n(p), g)$ and $(GLSym_n(n-p), g)$; in particular (\mathcal{P}_n, g) and $(GLSym_n(0), g)$ are isometric Riemannian manifolds.

Proof. a) Let $A \in GLSym_n(p)$ and $C \in GL_n$ such that $CAC^T = \Gamma_C(A) = J_p$. Since the restriction $\Gamma_C|_{GLSym_n(p)}$ is an isometry of $(GLSym_n(p), g)$, then $(GLSym_n(p), g)$ is homogeneous for any $p = 0, \dots, n$. Hence to prove that $(GLSym_n(p), g)$ is a Semi-Riemannian submanifold of (GL_n, g) , it suffices to verify that g_{J_p} is non-degenerate on $T_{J_p}(GLSym_n(p)) = Sym_n$ with the expected signature.

Since the set \mathcal{B} is clearly a basis of the vector space Sym_n , it suffices to compute g_{J_p} on the pairs of elements of \mathcal{B} .

For every $X = (x_{ij}), Y = (y_{ij}) \in Sym_n$, standard computations allow to get: $g_{J_p}(X, Y) = tr(J_p X J_p Y) = \sum_{i=1}^n x_{ii} y_{ii} + \sum_{1 \leq i < j \leq p} 2x_{ij} y_{ij} + \sum_{p+1 \leq i < j \leq n} 2x_{ij} y_{ij} - \sum_{1 \leq i \leq p < j \leq n} 2x_{ij} y_{ij}$.

This formula allows to conclude part (a) by direct computations.

b) The assertion follows by trivial checks. \square

2.2 Proposition. Fix $p \in \{0, \dots, n\}$. The inversion ϕ and the congruences Γ_C ($C \in GL_n$) are isometries of $(GLSym_n(p), g)$.

If $C \in GLSym_n(p)$, then the isometry $\phi_C = \Gamma_C \circ \phi$ is the symmetry of $(GLSym_n(p), g)$ fixing C ; therefore $(GLSym_n(p), g)$ is a symmetric Semi-Riemannian manifold.

Proof. The first part is trivial. For the second one it suffices to argue on the symmetry $\phi_C : X \mapsto CX^{-1}C^T = CX^{-1}C$ with respect to every $C \in GLSym_n$. \square

2.3 Proposition. a) For any $p = 0, \dots, n$, $(GLSym_n(p), g)$ is a totally geodesic submanifold of (GL_n, g) and its geodesics are precisely the curves of type $t \mapsto Ke^{tC}$ for every $K \in GLSym_n(p)$ and every $C = K^{-1}V$ with $V \in Sym_n$. In particular $(GLSym_n(p), g)$ is geodesically complete.

b) The Levi-Civita connection ∇ of $(GLSym_n(p), g)$ is

$$(\nabla_X \mathcal{Y})_K = (X(\mathcal{Y}))_K - \frac{1}{2}(X_K K^{-1} \mathcal{Y}_K + \mathcal{Y}_K K^{-1} X_K),$$

for every $K \in GLSym_n(p)$ and for all tangent vector fields X, \mathcal{Y} of class C^∞ on $GLSym_n(p)$;

the Riemann curvature tensor of type $(0, 4)$ of $(GLSym_n(p), g)$ is

$$R_{XYZW}(K) = \frac{1}{4} \text{tr}([K^{-1}X, K^{-1}Y][K^{-1}Z, K^{-1}W]),$$

for every $K \in GLSym_n(p)$ and every $X, Y, Z, W \in Sym_n = T_K(GLSym_n(p))$.

c) Let $\gamma = \gamma(t)$ (with t in a real interval \mathcal{J} containing 0) be a C^∞ -curve contained in $GLSym_n(p)$ and $\tau_{\gamma(t)} : T_{\gamma(0)}(GLSym_n(p)) \rightarrow T_{\gamma(t)}(GLSym_n(p))$ be the parallel transport along $\gamma(t)$ induced by the Levi-Civita connection of $(GLSym_n(p), g)$. Then $\tau_{\gamma(t)}$ agrees with the congruence $\Gamma_{F(t)^{-1}}$:

$$\tau_{\gamma(t)} = \Gamma_{F(t)^{-1}} \quad \forall t \in \mathcal{J},$$

where $F = F(t)$ is the unique (non-singular) solution, for every $t \in \mathcal{J}$, of the following matrixial system:

$$\begin{cases} \dot{F} = -\frac{1}{2}F\dot{\gamma}\gamma^{-1} \\ F(0) = I_n. \end{cases}$$

Proof. a) Remembering 1.3 (a), it suffices to check that $Ke^{tC} \in GLSym_n(p)$ for every $K \in GLSym_n(p)$, for every $C = K^{-1}V$ with $V \in Sym_n$ and for every $t \in \mathbb{R}$.

For, by standard properties of the exponential mapping: $(Ke^{tK^{-1}V})^T = e^{tVK^{-1}}K = Ke^{tK^{-1}V}K^{-1}K = Ke^{tK^{-1}V}$. So $Ke^{tK^{-1}V} \in GLSym_n(p)$.

b) It follows by 1.3 (b), since $(GLSym_n(p), g)$ is a totally geodesic submanifold of (GL_n, g) .

c) It is a standard fact that the system in the statement has a unique C^∞ -solution which is never singular (see for instance [1, Thm. 7.15, pp. 219–220]). By (b), the equation of the parallel transport is: $\dot{\mathcal{Y}} = \frac{1}{2}(\dot{\gamma}\gamma^{-1}\mathcal{Y} + \mathcal{Y}\gamma^{-1}\dot{\gamma})$. A direct computation shows that $\mathcal{Y}(t) = F(t)^{-1}WF(t)^{-T} = \Gamma_{F(t)^{-1}}(W)$ is its solution under the condition $\mathcal{Y}(0) = W$ and this allows to conclude. \square

2.4 Remark. Let $\gamma(t) = Ke^{tK^{-1}V} = \Gamma_{\exp(\frac{1}{2}tVK^{-1})}(K)$ be the geodesic of $(GLSym_n(p), g)$ such that $\gamma(0) = K \in GLSym_n(p)$ and $\dot{\gamma}(0) = V \in Sym_n$. In this case the parallel transport along $\gamma(t)$ becomes $\tau_{\gamma(t)} = \Gamma_{\exp(\frac{1}{2}tVK^{-1})}$ for every $t \in \mathbb{R}$, in accordance with the general theory (see for instance [15, Ch. XI, Thm. 3.2]).

2.5 Proposition. For every $p \in \{0, \dots, n\}$, the Ricci curvature of $(GLSym_n(p), g)$ is

$$\text{Ric}_Q(X, Z) = \frac{1}{4} \text{tr}(Q^{-1}X) \text{tr}(Q^{-1}Z) - \frac{n}{4} g_Q(X, Z)$$

for every $Q \in GLSym_n(p)$ and for every $X, Z \in T_Q(GLSym_n(p)) = Sym_n$ and the scalar curvature of $(GLSym_n(p), g)$ is

$$S = -\frac{(n-1)n(n+2)}{8}.$$

Proof. Fixed $p \in \{0, \dots, n\}$, we denote $\varepsilon_i = 1$, for $i = 1, \dots, p$, $\varepsilon_i = -1$, for $i = p + 1, \dots, n$, and $J = J_p$. If $X, Y \in \mathbb{R}^n$ are column vectors, we denote: $\langle X, Y \rangle = X^T J Y = \sum_{i=1}^n \varepsilon_i x_i y_i$, where $X = (x_1, \dots, x_n)^T$ and $Y = (y_1, \dots, y_n)^T$ (the *Minkowski product* of \mathbb{R}^n of signature $(p, n-p)$).

Given every matrix A , we denote by A^h its h -th column. If A is symmetric, A^h is the h -th row too. From 2.3 (b), by standard computation we get:

$$\begin{aligned} -4R_{XYZW} &= \\ &= \sum_{i,j=1}^n \varepsilon_i \varepsilon_j (\langle X^i, Y^j \rangle \langle Z^j, W^i \rangle + \langle Y^i, X^j \rangle \langle W^j, Z^i \rangle + \\ &\quad - \langle Y^i, X^j \rangle \langle Z^j, W^i \rangle - \langle X^i, Y^j \rangle \langle W^j, Z^i \rangle), \end{aligned}$$

for every $X, Y, Z, W \in T_J(GLSym_n(p)) = Sym_n$.

The basis \mathcal{B} of 2.1 can be listed also by $\{W_{(i,j)} = \frac{E^{(i,j)} + E^{(j,i)}}{\sqrt{2(1 + \delta_{ij})}} \mid 1 \leq i \leq j \leq n\}$ and so $g(W_{(i,j)}, W_{(i,j)}) = \varepsilon_i \varepsilon_j$ for every $1 \leq i \leq j \leq n$.

From the previous formula, for all symmetric matrices X, Z we get:

$$\begin{aligned} 4Ric_J(X, Z) &= -4 \sum_{1 \leq k \leq m \leq n} g(W_{(k,m)}, W_{(k,m)}) R_{XW_{(k,m)}ZW_{(k,m)}} = \\ &= 2 \sum_{1 \leq k \leq m \leq n} \sum_{i,j=1}^n \varepsilon_k \varepsilon_m \varepsilon_i \varepsilon_j (\langle X^i, W_{(k,m)}^j \rangle \langle Z^j, W_{(k,m)}^i \rangle + \\ &\quad - \langle X^j, W_{(k,m)}^i \rangle \langle Z^i, W_{(k,m)}^j \rangle). \end{aligned}$$

For every symmetric matrix $A = (a_{ij})$ we have $\langle A^i, W_{(k,m)}^j \rangle = \frac{a_{ik} \varepsilon_k \delta_{jm} + a_{im} \varepsilon_m \delta_{jk}}{\sqrt{2(1 + \delta_{km})}}$.

Hence, by standard computations, if $X = (x_{ij}), Z = (z_{ij}) \in Sym_n$ we get:

$$\begin{aligned} 4Ric_J(X, Z) &= \\ &= \sum_{1 \leq k \leq m \leq n} \frac{1}{1 + \delta_{km}} [\varepsilon_k \varepsilon_m x_{mk} z_{mk} + \varepsilon_k \varepsilon_m x_{kk} z_{mm} + \varepsilon_m \varepsilon_k x_{mm} z_{kk} + \varepsilon_m \varepsilon_k x_{km} z_{km} + \\ &\quad - \sum_{j=1}^n (\varepsilon_k \varepsilon_j x_{jk} z_{jk} + \varepsilon_k \varepsilon_j x_{jm} z_{jk} \delta_{mk} + \varepsilon_k \varepsilon_j x_{jk} z_{jm} \delta_{mk} + \varepsilon_m \varepsilon_j x_{jm} z_{jm})] = \\ &= 2 \sum_{k=1}^n x_{kk} z_{kk} + \sum_{1 \leq k < m \leq n} \varepsilon_k \varepsilon_m (x_{kk} z_{mm} + x_{mm} z_{kk} + 2x_{km} z_{km}) + \\ &\quad - 2 \sum_{k,j=1}^n \varepsilon_k \varepsilon_j x_{jk} z_{jk} - \sum_{1 \leq k < m \leq n} \varepsilon_k \sum_{j=1}^n \varepsilon_j x_{jk} z_{jk} - \sum_{1 \leq k < m \leq n} \varepsilon_m \sum_{j=1}^n \varepsilon_j x_{jm} z_{jm}. \end{aligned}$$

Note that: $tr(JX)tr(JZ) + g_J(X, Z) =$

$$= 2 \sum_{k=1}^n x_{kk} z_{kk} + \sum_{1 \leq k < m \leq n} \varepsilon_k \varepsilon_m (x_{kk} z_{mm} + x_{mm} z_{kk} + 2x_{km} z_{km}).$$

Moreover: $-2g_J(X, Z) = 2 \sum_{k,j=1}^n \varepsilon_k \varepsilon_j x_{jk} z_{jk}$.

Furthermore:

$$-(n-1)g_J(X, Z) = -\sum_{1 \leq k < m \leq n} \varepsilon_k \sum_{j=1}^n \varepsilon_j x_{jk} z_{jk} - \sum_{1 \leq k < m \leq n} \varepsilon_m \sum_{j=1}^n \varepsilon_j x_{jm} z_{jm}.$$

Comparing this with the expression of $4Ric_J(X, Z)$, we get that:

$$\begin{aligned} 4Ric_J(X, Z) &= tr(JX)tr(JZ) + g_J(X, Z) - 2g_J(X, Z) - (n-1)g_J(X, Z) = \\ &= tr(JX)tr(JZ) - ng_J(X, Z). \end{aligned}$$

Hence:

$$Ric_J(X, Z) = \frac{1}{4} tr(JX)tr(JZ) - \frac{n}{4} g_J(X, Z), \text{ for every } X, Z \in T_J(GLSym_n(p)) = Sym_n.$$

Now let Q be a generic matrix of $GLSym_n(p)$. We know that there exists a non-singular C such that $\Gamma_C(Q) = CQC^T = J$, i.e. $JC = C^{-T}Q^{-1}$. Since Γ_C is an isometry of $(GSym_n(p), g)$, it preserves Ric , i.e. for any $X, Z \in Sym_n$:

$$\begin{aligned} Ric_Q(X, Z) &= Ric_{\Gamma_C(Q)}(\Gamma_C(X), \Gamma_C(Z)) = Ric_J(CXC^T, CZC^T) = \\ &= \frac{1}{4}tr(JCXC^T)tr(JCZC^T) - \frac{n}{4}tr(JCXC^TJCZC^T) = \frac{1}{4}tr(Q^{-1}X)tr(Q^{-1}Z) - \frac{n}{4}g_Q(X, Z) \end{aligned}$$

and this concludes the first part.

The scalar curvature S of $(GLSym_n(p), g)$ is constant, because this manifold is homogeneous. So it suffices to compute S at the point J . Now $J \in T_J(GLSym_n(p)) = Sym_n$ is a space-like vector (indeed $g_J(J, J) = n$), hence we have:

$$Sym_n = T_J(GLSym_n(p)) = Span(J) \oplus (J)^\perp, \text{ where } (J)^\perp = \{V \in Sym_n : g_J(J, V) = 0\}.$$

Let V_1, \dots, V_d with $d = \frac{n(n+1)}{2} - 1$ be an orthonormal basis of $(J)^\perp$. We have:

$$S = g_J\left(\frac{J}{\sqrt{n}}, \frac{J}{\sqrt{n}}\right) Ric_J\left(\frac{J}{\sqrt{n}}, \frac{J}{\sqrt{n}}\right) + \sum_{i=1}^d g_J(V_i, V_i) Ric_J(V_i, V_i).$$

Now from the expression of Ric , the latter is equal to:

$$\begin{aligned} \frac{1}{4} \left[tr\left(\frac{J^2}{\sqrt{n}}\right) \right]^2 - \frac{n}{4} tr\left(\frac{J^4}{n}\right) + \sum_{i=1}^d \frac{1}{4} g_J(V_i, V_i) [tr(JV_i)]^2 - \frac{n}{4} \sum_{i=1}^d g_J(V_i, V_i) g_J(V_i, V_i) = \\ -\frac{n}{4}d, \text{ since } tr(JV_i) = g_J(J, V_i) = 0 \text{ for every } i = 0, \dots, d. \end{aligned}$$

$$\text{Hence: } S = -\frac{n}{4} \left(\frac{n(n+1)}{2} - 1 \right) = -\frac{(n-1)n(n+2)}{8}. \quad \square$$

2.6 Proposition. For every $p = 0, \dots, n$ let us consider the set $SLSym_n(p)$.

a) $(SLSym_n(p), g)$ is a homogeneous Semi-Riemannian submanifold of $(GLSym_n(p), g)$

with signature $\left(\frac{n(n+1)}{2} - p(n-p) - 1, p(n-p)\right)$.

b) $(SLSym_n(p), g)$ is a totally geodesic submanifold of $(GLSym_n(p), g)$, whose geodesics are the curves $t \mapsto Kexp(tK^{-1}V)$ for every $K \in SLSym_n(p)$ and every $V \in Sym_n$ with $tr(K^{-1}V) = 0$. In particular $(SLSym_n(p), g)$ is geodesically complete.

c) The inversion ϕ and the congruences Γ_C with $\det(C) = \pm 1$ are isometries of the symmetric manifold $(SLSym_n(p), g)$ and, for every Q , ϕ_Q is the symmetry fixing Q .

d) $(SLSym_n(p), g)$ is an Einstein manifold with Ricci tensor $Ric = -\frac{n}{4}g$ and with scalar curvature $S = -\frac{(n-1)n(n+2)}{8}$.

Proof. a) Note that $SLSym_n(p) = GLSym_n(p) \cap SL_n((-1)^{n-p})$, where $SL_n((-1)^{n-p}) = \{A \in GL_n : \det(A) = (-1)^{n-p}\}$ is a submanifold of GL_n of codimension 1 such that, for every $Q \in SL_n((-1)^{n-p})$, by the well-known Jacobi's formula, we have:

$$T_Q(SL_n((-1)^{n-p})) = \{V \in M_n : tr(Q^{-1}V) = 0\}.$$

Hence: $T_Q(GL_n) = M_n$ is the sum of its vector subspaces $T_Q(GLSym_n(p)) = Sym_n$ and $T_Q(SL_n((-1)^{n-p}))$, since $Q \in T_Q(GLSym_n(p)) \setminus T_Q(SL_n((-1)^{n-p}))$.

Thus $GLSym_n(p)$ and $SL_n((-1)^{n-p})$ intersect transversally and therefore $SLSym_n(p) = GLSym_n(p) \cap SL_n((-1)^{n-p})$ is a submanifold of $GLSym_n(p)$ of dimension $\frac{n(n+1)}{2} - 1$ (see [12, Thm. 3.3, p. 22]).

Of course $T_Q(SLSym_n(p)) = \{V \in Sym_n : tr(Q^{-1}V) = 0\}$ for every $Q \in SLSym_n(p)$. With the same notations as in the proof of 2.5, for $J = J_p$, we have: $T_J(SLSym_n(p)) = (J)^\perp = \{V \in Sym_n : g_J(J, V) = 0\}$. Since J is a space-like vector of $T_J(GLSym_n(p))$, the restriction of g_J to $T_J(SLSym_n(p))$ is non-degenerate with signature

$(\frac{n(n+1)}{2} - p(n-p) - 1, p(n-p))$. Now if $Q \in SLSym_n(p)$, there exists a non-singular matrix C such that $\Gamma_C(Q) = CQC^T = J$. Hence $det(C) = \pm 1$ and Γ_C is an isometry of $(GLSym_n(p), g)$, mapping $SLSym_n(p)$ onto itself and Q to J . We conclude that g_J and g_Q have the same signature, for every Q .

b) By 2.3 (a), it suffices to check that $Qe^{tC} \in SLSym_n(p)$ for every $Q \in SLSym_n(p)$, for every $C = Q^{-1}V$ with $V \in T_Q(SLSym_n(p))$ and for every $t \in \mathbb{R}$.

For, it suffices to compute $det(Qe^{tC})$ via the fact that $tr(Q^{-1}V) = 0$.

c) The mappings in the statement are clearly isometries. If $Q \in SLSym_n(p)$, then $\varphi_Q = \Gamma_Q \circ \varphi$ is the expected symmetry of $(SLSym_n(p), g)$, fixing Q .

d) If $Q \in SLSym_n(p)$ then $N = N_Q = \frac{Q}{\sqrt{n}} \in T_Q(GLSym_n(p)) = Sym_n$ is a space-like unit vector and $T_Q(SLSym_n(p)) = (N)^\perp$. Since $T_Q(GLSym_n(p)) = Span(N) \oplus (N)^\perp$, if V_1, \dots, V_d (with $d = \frac{n(n+1)}{2} - 1$) is an orthonormal basis of $T_Q(SLSym_n(p))$, then N, V_1, \dots, V_d is an orthonormal basis of $T_Q(GLSym_n(p))$.

Hence, if $X, Z \in T_Q(GLSym_n(p))$ and Ric_Q is the Ricci tensor at Q of $(GLSym_n(p), g)$, we have: $Ric_Q(X, Z) = -(R_{XNZN} + \sum_{i=1}^d g_Q(V_i, V_i)R_{XV_iZV_i}) = -\sum_{i=1}^d g_Q(V_i, V_i)R_{XV_iZV_i}$, being $R_{XNZN} = 0$, by 2.3 (b).

By part (b), the Riemann tensor of $(SLSym_n(p), g)$ is the restriction to $SLSym_n(p)$ of the Riemann tensor of $(GLSym_n(p), g)$. From the previous expression of $Ric_Q(X, Z)$, we deduce that the restriction to $T_Q(SLSym_n(p))$ of the Ricci tensor of $(GLSym_n(p), g)$ coincides with the Ricci tensor of $(SLSym_n(p), g)$ at Q . Hence, if $X, Z \in T_Q(SLSym_n(p))$, by part (a), the Ricci tensor of $(SLSym_n(p), g)$ at Q is

$$Ric_Q(X, Z) = \frac{1}{4}\{tr(Q^{-1}X)tr(Q^{-1}Z) - n g_Q(X, Z)\} = -\frac{n}{4}g_Q(X, Z).$$

Hence $(SLSym_n(p), g)$ is an Einstein manifold with $Ric = -\frac{n}{4}g$ and with

$$S = -\frac{n}{4}dim(SLSym_n(p)) = -\frac{(n-1)n(n+2)}{8}. \quad \square$$

2.7 Proposition. For every $p = 0, \dots, n$, $(GLSym_n(p), g)$ is isometric to the Semi-Riemannian product manifold $(SLSym_n(p) \times \mathbb{R}, g \times h)$, with h euclidean metric on \mathbb{R} .

Proof. The mapping $F : (SLSym_n(p) \times \mathbb{R}, g \times h) \rightarrow (GLSym_n(p), g)$, given by

$$F(Q, x) = e^{\frac{x}{\sqrt{n}}}Q, \text{ is invertible with inverse } F^{-1}(P) = \left(\frac{P}{\sqrt{|\det(P)|}}, \frac{\ln(|\det(P)|)}{\sqrt{n}}\right). \text{ We}$$

can easily check that F and F^{-1} are isometries (see also [6, Thm. 4.2]). \square

3. The Riemannian manifold (\mathcal{P}_n, g)

3.1 Proposition. *a) (\mathcal{P}_n, g) and $(SL\mathcal{P}_n, g)$ are simply connected, symmetric Riemannian manifolds with non-positive sectional curvature; (\mathcal{P}_n, g) is isometric to the Riemannian product manifold $SL\mathcal{P}_n \times \mathbb{R}$ and $SL\mathcal{P}_n$ cannot be expressed as a non-trivial (metric) product.*

b) The Levi-Civita connection ∇ of (\mathcal{P}_n, g) is

$$(\nabla_X \mathcal{Y})_K = (X(\mathcal{Y}))_K - \frac{1}{2}(\mathcal{X}_K K^{-1} \mathcal{Y}_K + \mathcal{Y}_K K^{-1} \mathcal{X}_K),$$

for every $K \in \mathcal{P}_n$ and for all tangent vector fields X, \mathcal{Y} of class C^∞ on \mathcal{P}_n ;

furthermore, if $\gamma(t)$, $\tau_{\gamma(t)}$, $F(t)$ and \mathcal{J} are as in 2.3 (c), then the parallel transport, $\tau_{\gamma(t)}$, along $\gamma(t)$ induced by the Levi-Civita connection of (\mathcal{P}_n, g) is given by

$$\tau_{\gamma(t)} = \Gamma_{F(t)}^{-1} \quad \forall t \in \mathcal{J}.$$

c) The Ricci curvature of (\mathcal{P}_n, g) at any point Q is negative semi-definite and $Ric_Q(X, X) = 0$ if and only if $X = \lambda Q$, for some $\lambda \in \mathbb{R}$; the scalar curvature is $-\frac{(n-1)n(n+2)}{8}$.

d) $(SL\mathcal{P}_n, g)$ is an Einstein manifold with Ricci tensor $Ric = -\frac{n}{4}g$ and with scalar curvature $S = -\frac{(n-1)n(n+2)}{8}$.

Proof. a) These results are well-known: we refer for instance to [25, Thm. 2.1] for the curvature and to [5, Prop. 10.34, Lemma 10.52, Prop. 10.53] for the remaining facts.

However, except for the sectional curvature and for the irreducibility of $SL\mathcal{P}_n$, these results are also direct consequences of the more general facts proved in the previous Section. For completeness we get also the remaining assertions.

For the curvature, by homogeneity, it suffices to compute the sectional curvature at point I_n , where, by 2.3, we have $R_{XYXY} = \frac{1}{4}tr([X, Y]^2)$. Since $[X, Y]$ is skew-symmetric, $tr([X, Y]^2)$ is the opposite of the sum of the squares of the entries of the matrix $[X, Y]$ (i.e. the opposite of the *Frobenius norm* of $[X, Y]$) and so the sectional curvature is non-positive.

Finally, it is known that $SL\mathcal{P}_n \simeq SL_n/SO_n$ is an irreducible symmetric space (see for instance [2, Table 3 p. 315]), hence $(SL\mathcal{P}_n \times \mathbb{R}, g \times h)$ is the *de Rham decomposition* of the simply connected complete Riemannian manifold (\mathcal{P}_n, g) (see for instance [14, Ch. IV, Thm. 6.2]).

b) It is a particular case of Proposition 2.3 (b) and (c). The Levi-Civita connection is also in [25, p. 214] and in [19, p. 176].

c) The assertion about the scalar curvature is in 2.6 (d) (and also quoted in [19, p. 176] with opposite sign).

Since Q is symmetric positive definite, there exists a non-singular matrix C such that $Q = CC^T = \Gamma_C(I_n)$. Let $X \in T_Q(\mathcal{P}_n) = \text{Sym}_n$, then by proposition 2.5:

$$4\text{Ric}_Q(X, X) = [\text{tr}(Q^{-1}X)]^2 - n g_Q(X, X) = [\text{tr}(Y)]^2 - n \text{tr}(Y^2),$$

where $Y = C^{-1}XC^{-T}$ is a symmetric matrix. The statement follows from the following:

Claim.

For every symmetric real matrix Y of order n , we have $[\text{tr}(Y)]^2 \leq n \text{tr}(Y^2)$ with equality if and only if $Y = \lambda I_n$ for some $\lambda \in \mathbb{R}$.

Indeed, if $\lambda_1, \dots, \lambda_n$ are the (possibly repeated) real eigenvalues of Y , then

$$[\text{tr}(Y)]^2 = \sum_{i,j=1}^n \lambda_i \lambda_j \leq \sum_{i,j=1}^n \frac{\lambda_i^2 + \lambda_j^2}{2} = n \sum_{i=1}^n \lambda_i^2 = n \text{tr}(Y^2)$$

and the equality holds if and only if $\lambda_i = \lambda_j$ for every i, j , i.e. if and only if $Y = \lambda I_n$ for some $\lambda \in \mathbb{R}$, being Y diagonalizable.

d) It follows directly from 2.6 (d), in accordance with the general theory (see for instance [2, Note 10.83, p. 298]). \square

3.2 Remarks. a) For every point $Q \in \mathcal{P}_n$, the corresponding symmetry ϕ_Q has Q as unique fixed point.

Indeed, by homogeneity, it is enough to check this for the case $Q = I_n \in \mathcal{P}_n$. We have $\phi_{I_n} = \phi$ and so $X \in \mathcal{P}_n$ is a fixed point of ϕ if and only if X is orthogonal too, therefore $X = I_n$.

b) $(SL\mathcal{P}_2, g)$ is isometric to the hyperbolic plane \mathcal{H}_2 (endowed with Riemannian metric having curvature $-\frac{1}{2}$).

Indeed $(SL\mathcal{P}_2, g)$ is a complete, simply connected, homogeneous, Riemannian surface and, therefore, its curvature is constant and equal to $\frac{S}{2} = -\frac{1}{2}$ (see for instance [14, Ch. VI, Thm. 7.10]).

Hence \mathcal{P}_2 is isometric to the Riemannian product $\mathcal{H}_2 \times \mathbb{R}$.

c) As in 2.4, if $\gamma(t) = Ke^{tK^{-1}V}$ is the geodesic of (\mathcal{P}_n, g) such that $\gamma(0) = K \in \mathcal{P}_n$ and $\dot{\gamma}(0) = V \in \text{Sym}_n$, then: $\tau_{\gamma(t)} = \Gamma_{\exp(\frac{1}{2}tVK^{-1})}$ for every $t \in \mathbb{R}$.

3.3 Remark-Definition. Let M be an $n \times n$ real matrix, diagonalizable over \mathbb{R} with (possibly repeated) eigenvalues $\lambda_1, \dots, \lambda_n$, all strictly positive, and $G \in GL_n$ be a matrix such that $M = G^{-1} \text{diag}(\lambda_1, \dots, \lambda_n)G$.

We denote by $\text{LOG}(M)$ the matrix $G^{-1} \text{diag}(\ln(\lambda_1), \dots, \ln(\lambda_n))G$.

$\text{LOG}(M)$ is the unique solution of the equation $\exp(X) = M$ among the $n \times n$ real matrices, which are diagonalizable over \mathbb{R} ; the proof is contained in [11, Thm. 1.31]. $\text{LOG}(M)$ is said to be the *principal logarithm* of M . Therefore, for every $r \in \mathbb{R}$, it is possible to define the r -th power of M as $M^r = \exp(r \text{LOG}(M))$.

3.4 Proposition. Let $A, B \in \mathcal{P}_n$.

a) $\gamma(t) = A \exp(t \text{LOG}(A^{-1}B)) = A(A^{-1}B)^t$ is the unique geodesic arc $\gamma(t) : [0, 1] \rightarrow (\mathcal{P}_n, g)$ such that $\gamma(0) = A$ and $\gamma(1) = B$.

b) The distance $d(A, B)$ between the matrices $A, B \in \mathcal{P}_n$, induced by the trace metric, is

$$d(A, B) = (\sum_{i=1}^n (\ln \mu_i)^2)^{1/2}$$

where μ_1, \dots, μ_n are the (possibly repeated) eigenvalues of $A^{-1}B$.

Proof. These results are known (see [4, § 2], [3, Ch. 6 § 1] and [19, § 3.5, § 3.6]). We shortly prove them, by using the arguments previously developed.

The classical Theorem of Cartan-Hadamard ([22, Ch. 10 Thm. 22]) implies that there is a unique geodesic arc as in (a) and by Hopf-Rinow ([22, Ch. 5 Thm. 21]) its length gives $d(A, B)$.

By simultaneous diagonalization, there is a non-singular matrix C such that $\Gamma_C(A) = CAC^T = I_n$ (i.e. $A = C^{-1}C^{-T}$) and $\Gamma_C(B) = D := \text{diag}(\mu_1, \dots, \mu_n)$, where μ_1, \dots, μ_n are the (necessarily positive) eigenvalues of $A^{-1}B$, which is diagonalizable over \mathbb{R} .

Now $\text{LOG}(D) = \text{diag}(\ln(\mu_1), \dots, \ln(\mu_n))$ and so by 2.3, the unique geodesic arc joining I_n and D is $\beta(t) = \exp(t \text{LOG}(D))$, $t \in [0, 1]$.

Since Γ_C is an isometry, we get: $d(A, B) = d(I_n, D) = \text{length}(\beta) = [g_{I_n}(\dot{\beta}(0), \dot{\beta}(0))]^{1/2} = [g_{I_n}(\text{LOG}(D), \text{LOG}(D))]^{1/2} = (\sum_{i=1}^n (\ln \mu_i)^2)^{1/2}$.

Now the unique geodesic arc joining A and B is

$$\gamma(t) = \Gamma_{C^{-1}}(\beta(t)) = C^{-1} \exp(t \text{LOG}(D)) C^{-T} = A \exp(t C^T \text{LOG}(D) C^{-T}).$$

Note that, by 3.3, $C^T \text{LOG}(D) C^{-T} = \text{LOG}(A^{-1}B)$. This allows to conclude. \square

3.5 Remark. The description of the full group of isometries of (\mathcal{P}_n, g) is in §4. For now we recall that inversion and congruences are isometries of (\mathcal{P}_n, g) , so GL_n acts transitively by congruences on the Riemannian manifold (\mathcal{P}_n, g) (remember 1.2 and 2.2).

Moreover it is possible to prove that for every pair $A, B \in \mathcal{P}_n$ there is a unique matrix $X \in \mathcal{P}_n$ such that $\Gamma_X(A) = B$ and that X is the *geometric mean* of the matrices A^{-1} and B , i.e. the *midpoint* of the unique geodesic joining A^{-1} and B in the manifold (\mathcal{P}_n, g) (see for instance [3, p. 11, pp. 106–107, p. 206]). In particular the homogeneity of (\mathcal{P}_n, g) can be obtained by means of congruences associated to positive definite matrices (see also [16, Ch. XII, Lemma 3.2]).

3.6 Proposition. Fixed a matrix $U \in O_n$, we denote: $\hat{\mathcal{L}}_U = \{UQ \in GL_n : Q \in \mathcal{P}_n\}$ and $\hat{\mathcal{R}}_U = \{QU \in GL_n : Q \in \mathcal{P}_n\}$. Then $\hat{\mathcal{L}}_U = \hat{\mathcal{R}}_U$ and $GL_n = \bigcup_{U \in O_n} \hat{\mathcal{L}}_U$ is a foliation of (GL_n, g) , whose leaves are isometric to (\mathcal{P}_n, g) and totally geodesic in (GL_n, g) .

Proof. From the *polar decomposition* (see for instance [13, Thm. 7.3.1]), for every matrix $A \in GL_n$ there is a unique $U \in O_n$ and there are unique $Q, Q' \in \mathcal{P}_n$ such that $A = UQ = Q'U$ (so: $Q' = UQU^T$). This gives that each $A \in GL_n$ belongs to a unique $\hat{\mathcal{L}}_U$ and gives also the equality $\hat{\mathcal{L}}_U = \hat{\mathcal{R}}_U$. We get the last part of the statement, because, by [6, Prop. 1.2], left translations are isometries of (GL_n, g) (remember also 2.3 (a)). \square

3.7 Remark. The roles of O_n and of \mathcal{P}_n in the previous Proposition are mutually interchangeable: indeed, in [7, Prop. 4.5], we proved that GL_n has analogous foliations:

$GL_n = \bigcup_{Q \in \mathcal{P}_n} \mathcal{L}_Q = \bigcup_{Q \in \mathcal{P}_n} \mathcal{R}_Q$ whose leaves are all isometric to (O_n, g) and totally geodesic in (GL_n, g) .

4. The group of isometries of (\mathcal{P}_n, g)

Next Lemma and next Proposition are known, but we state and shortly prove them for convenience and completeness.

4.1 Lemma. *Let \mathbf{G} be a connected Lie group and $\tilde{\mathbf{G}}$ be its universal covering group. Then $Out(\mathbf{G})$ is isomorphic to a subgroup of $Out(\tilde{\mathbf{G}})$.*

Proof. For every $\alpha \in Aut(\mathbf{G})$ we denote by $\tilde{\alpha} : \tilde{\mathbf{G}} \rightarrow \tilde{\mathbf{G}}$ the unique lift of α such that $\tilde{\alpha}(1_{\tilde{\mathbf{G}}}) = 1_{\tilde{\mathbf{G}}}$; then $\tilde{\alpha}$ is in $Aut(\tilde{\mathbf{G}})$ and the mapping which associates to α the equivalence class of $\tilde{\alpha}$ in $Out(\tilde{\mathbf{G}})$ is a group homomorphism with kernel $Inn(\mathbf{G})$. \square

4.2 Proposition. *$Aut(SO_n) \simeq SO_n$ if n is odd and $Aut(SO_n/\{\pm I_n\}) \simeq O_n/\{\pm I_n\}$ if $n \neq 8$ is even, where the groups on the right act by conjugation.*

Proof. The following arguments need some facts recalled in 1.1.

If n is odd, the result follows from the previous Lemma and from the fact that $Out(Spin_n)$ is trivial.

If n is even and different from 2, then $Spin_n$ is also the universal covering of $SO_n/\{\pm I_n\}$. If $n \neq 8$ too, by the previous Lemma, $Out(SO_n/\{\pm I_n\})$ has at most two elements. Since any conjugation by elements of $O_n \setminus SO_n$ is an outer automorphism of $SO_n/\{\pm I_n\}$, then $Out(SO_n/\{\pm I_n\}) \simeq \mathbb{Z}_2$ and this gives the assertion.

If $n = 2$, then $SO_2/\{\pm I_2\} \simeq S^1$ (the circle), $Aut(S^1) = Out(S^1) \simeq \mathbb{Z}_2$ (with elements the identity and the complex inversion) and so $Out(SO_2/\{\pm I_2\}) \simeq \mathbb{Z}_2$ and this allows to conclude as above. \square

Theorem 4.1. *A mapping $G : (SL\mathcal{P}_n, g) \rightarrow (SL\mathcal{P}_n, g)$ is an isometry if and only if there exists a matrix $X \in GL_n$ with $\det(X) = \pm 1$ such that (with the notations of 1.1)*

$$G = \Gamma_X \text{ or } G = \Gamma_X \circ \phi.$$

Moreover G fixes I_n if and only if the matrix X belongs to O_n .

Proof. The mappings Γ_X and $\Gamma_X \circ \phi$ with $\det(X) = \pm 1$ are isometries by 2.6 (c).

For the converse, up to congruences with matrices of determinant ± 1 , we can assume that G fixes I_n . Indeed $G(I_n) \in SL\mathcal{P}_n$ and so $G(I_n) = BB^T$ for some $B \in GL_n$ with $\det(B) = \pm 1$. Therefore $(\Gamma_{B^{-1}} \circ G)(I_n) = I_n$.

Let \mathcal{J} be the group of isometries of $(SL\mathcal{P}_n, g)$, \mathcal{J}_n be the corresponding subgroup of isotropy at I_n and $\mathcal{J}^0, \mathcal{J}_n^0$ be their connected components containing the identity.

Since $(SL\mathcal{P}_n, g)$ is homogeneous Riemannian (remember 3.1), we have: $SL\mathcal{P}_n \simeq \mathcal{J}/\mathcal{J}_n$. Remembering that $SL\mathcal{P}_n \simeq SL_n/SO_n$ (1.2 (b) and (c)), from [10, Ch. V Th. 4.1 (i)], we get that $\mathcal{J}^0 \simeq SL_n$ if n is odd and $\mathcal{J}^0 \simeq SL_n/\{\pm I_n\}$ if n is even.

Indeed it is well-known that SL_n is a connected simple Lie group and the actions of SL_n (if n is odd) and of $SL_n/\{\pm I_n\}$ (if n is even) are both effective.

From this we get that $\dim(\mathcal{J}) = \dim(\mathcal{J}^0) = \dim(SL_n)$ and therefore $\dim(SO_n) = \dim(\mathcal{J}_{I_n}) = \dim(\mathcal{J}_{I_n}^0)$.

Let us consider the representation $\rho : O_n \rightarrow \text{Aut}(\text{Sym}_n^0)$ defined by $\rho(X)(A) = XAX^T$. Arguing on the matrices $E^{(i,j)} + E^{(j,i)}$ and $E^{(i,i)} - E^{(j,j)}$ for every $i \neq j$, we get that $\text{Ker}(\rho) = \{\pm I_n\}$. Let us consider also the representation $d : \mathcal{J}_{I_n} \rightarrow \text{Aut}(\text{Sym}_n^0)$ defined by the differential at I_n of every element of \mathcal{J}_{I_n} (remember that $T_{I_n}(SL\mathcal{P}_n) = \text{Sym}_n^0$). By [22, Ch. 3 Prop. 62], d is a faithful representation and so: $d(\mathcal{J}_{I_n}^0) = (d(\mathcal{J}_{I_n}))^0$ (the component of $d(\mathcal{J}_{I_n})$ containing the identity).

Since congruences by orthogonal matrices are linear isometries fixing I_n , we get the inclusion $\rho(SO_n) \subseteq (d(\mathcal{J}_{I_n}))^0$. Since these manifolds have the same dimension and are connected, by *theorem of invariance of domain*, we deduce that $\rho(SO_n) = (d(\mathcal{J}_{I_n}))^0$.

For a fixed $G \in \mathcal{J}_{I_n}$, the previous equality gives: $dG\rho(SO_n)dG^{-1} = \rho(SO_n)$. Hence there exists a unique automorphism α of $\rho(SO_n)$ such that:

$$(*) \quad dG \circ \rho(X) \circ dG^{-1} = \alpha(\rho(X)) \text{ for every } X \in SO_n.$$

Claim. There is $Y \in O_n$ such that $\alpha(\rho(X)) = \rho(Y) \circ \rho(X) \circ \rho(Y)^{-1}$ for every $X \in SO_n$ and so, by (*), we get $(\rho(Y)^{-1} \circ dG) \circ \rho(X) = \rho(X) \circ (\rho(Y)^{-1} \circ dG)$ for every $X \in SO_n$.

Indeed $\rho(SO_n) \simeq SO_n$ if n is odd and $\rho(SO_n) \simeq SO_n/\{\pm I_n\}$ if n is even.

Hence, when $n \neq 8$, the claim follows by 4.2.

The case $n = 8$ needs different arguments.

First of all, we note that $F = \rho \circ \pi_8 : Spin_8 \rightarrow \rho(SO_8)$ is the universal covering of $\rho(SO_8)$; so the automorphism $\alpha \in \text{Aut}(\rho(SO_8))$ can be lifted to a unique $\tilde{\alpha} \in \text{Aut}(Spin_8)$ such that $F \circ \tilde{\alpha} = \alpha \circ F$.

As recalled in 1.1, $\text{Out}(Spin_8)$ is isomorphic to the dihedral group \mathbf{Dih}_3 and therefore it is the group of order 6 generated by elements δ and γ of order 2 and 3 respectively (see also [8] and [18] for further details). In particular we can assume that δ is the equivalence class of the lifting $\tilde{\tau}_H$ of the conjugation τ_H in SO_8 associated to a fixed matrix $H \in O_8 \setminus SO_8$ and γ is the equivalence class of an automorphism $\tilde{\gamma}$ of $Spin_8$ having the unit $1 \in Spin_8$ as unique fixed point in the fiber $\text{ker}(F)$ (see for instance [18, Ch. 1 §8]).

Therefore, up to inner automorphisms, we can assume that $\tilde{\alpha} = \tilde{\tau}_H^k \circ \tilde{\gamma}^p$ with $k = 0, 1$ and $p = 0, 1, 2$.

We prove that the unique admissible possibilities are $k = 0, 1$ and $p = 0$.

From (*) above, we deduce that $dG \circ F(Z) \circ dG^{-1} = \alpha(F(Z)) = F(\tilde{\alpha}(Z))$ for every $Z \in Spin_8$ (up to inner automorphisms); this implies that F and $F \circ \tilde{\alpha}$ are equivalent representations of $Spin_8$.

The cases $k = 0$ and $p = 1, 2$ are impossible because F , $F \circ \tilde{\gamma}$ and $F \circ \tilde{\gamma}^2$ are non-equivalent.

Indeed, by standard facts from Lie group representation theory (and with the help of the package *LiE* [26]), we get that the representations π_8 , $\pi_8 \circ \tilde{\gamma}$ and $\pi_8 \circ \tilde{\gamma}^2$ correspond

to maximal weights $(1, 0, 0, 0)$, $(0, 0, 1, 0)$ and $(0, 0, 0, 1)$ respectively, while their trace-free second symmetric powers are F , $F \circ \tilde{\gamma}$ and $F \circ \tilde{\gamma}^2$ and correspond to $(2, 0, 0, 0)$, $(0, 0, 2, 0)$ and $(0, 0, 0, 2)$ respectively and so they are mutually non-equivalent.

If $k = 1$ and $p = 1, 2$, since $\tilde{\tau}_H$ is the lifting of the automorphism τ_H of SO_8 , we can argue as above with $\rho(H)^{-1} \circ dG$ instead of dG . Therefore, up to inner automorphisms, $\tilde{\alpha} = \tilde{\tau}_H^k$ with $k = 0, 1$ and this allows to conclude the proof of the Claim.

Since the action of $\rho(SO_n)$ on Sym_n^0 is irreducible (see for instance [15, Ch. XI Prop. 7.4(1)] and [9, Prop. 4.5(1)]), by [14, App. 5 Lemma 1], we obtain: $\rho(Y)^{-1} \circ dG = aId + bJ$ (Id is the identity of Sym_n^0),

where $a, b \in \mathbb{R}$, $J^2 = -Id$. In particular, if $b \neq 0$, then $\dim(Sym_n^0)$ is even.

Now, if b would be non-zero, then the complexification of Sym_n^0 should have the eigenspaces of the complexification of J as invariant subspaces with respect to the complexification of the representation; so the complexification of the representation of $\rho(SO_n)$ would be reducible, while it is actually irreducible (see [9, Prop. 4.5(1), Prop. 4.6]).

Hence $b = 0$ and $\rho(Y)^{-1} \circ dG = aId$ and so, being an isometry, we get: $a = \pm 1$, i.e. $dG = \pm \rho(Y)$. In case of $a = 1$, then $G(A) = YAY^T = \Gamma_Y(A)$, while in case of $a = -1$, then $G(A) = YA^{-1}Y^T = \Gamma_Y(A^{-1}) = (\Gamma_Y \circ \phi)(A)$ (remember for instance [22, Ch. 3 Prop. 62]). This concludes the proof. \square

4.3 Remark. When $n = 2$, the inversion ϕ in $SL\mathcal{P}_2$ is the congruence associated to the rotation matrix $W = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$ and $I(SL\mathcal{P}_2, g)$ consists only of the indicated congruences and has two connected components, with $I^0(SL\mathcal{P}_2, g)$ isomorphic to $SL_2/\{\pm I_2\}$. If $n \geq 3$, then ϕ is not a congruence and the possibilities of the previous Theorem are mutually exclusive, moreover we get that:

- if n is odd, then $I(SL\mathcal{P}_n, g)$ is a Lie group with two connected components, with $I^0(SL\mathcal{P}_n, g)$ isomorphic to SL_n ;
- if $n \neq 2$ is even, then $I(SL\mathcal{P}_n, g)$ is a Lie group with four connected components, with $I^0(SL\mathcal{P}_n, g)$ isomorphic to $SL_n/\{\pm I_n\}$.

4.4 Proposition. A mapping $L : (\mathcal{P}_n, g) \rightarrow (\mathcal{P}_n, g)$ is an isometry if and only if there exist an isometry $G : (SL\mathcal{P}_n, g) \rightarrow (SL\mathcal{P}_n, g)$ and a matrix $B \in GL_n$ such that

$$L(A) = (\det A)^{1/n} (\Gamma_B \circ G) \left(\frac{A}{(\det A)^{1/n}} \right) \text{ for every } A \in \mathcal{P}_n \text{ or}$$

$$L(A) = (\det A)^{-1/n} (\Gamma_B \circ G) \left(\frac{A}{(\det A)^{1/n}} \right) \text{ for every } A \in \mathcal{P}_n.$$

Proof. By 3.1 (a) and 2.7, we have the isometry $F : SL\mathcal{P}_n \times \mathbb{R} \rightarrow \mathcal{P}_n$ given by

$$F(Q, x) = e^{\frac{x}{\sqrt{n}}} Q \text{ and } F^{-1}(P) = \left(\frac{P}{\sqrt[n]{\det(P)}}, \frac{\ln(\det(P))}{\sqrt{n}} \right). \text{ Hence } L : \mathcal{P}_n \rightarrow \mathcal{P}_n \text{ is an}$$

isometry if and only if $\bar{L} = F^{-1} \circ L \circ F$ is an isometry of $SL\mathcal{P}_n \times \mathbb{R}$.

Let \bar{L}^\pm be the isometries of $(SL\mathcal{P}_n \times \mathbb{R}, g \times h)$ defined by $\bar{L}^\pm(Q, x) = (G(Q), \pm x + b)$ where G is a fixed isometry of $(SL\mathcal{P}_n, g)$ and b is a fixed real number. Then $L^\pm =$

$F \circ \bar{L}^\pm \circ F^{-1}$ are isometries of (\mathcal{P}_n, g) . After denoting $B = e^{b/(2\sqrt{n})} I_n$, by standard computations, we get: $L^\pm(A) = \det(A)^{\pm 1/n} (\Gamma_B \circ G) \left(\frac{A}{(\det A)^{1/n}} \right)$.

For the converse, let L be an isometry of (\mathcal{P}_n, g) . Since $L(I_n) \in \mathcal{P}_n$, there exists $B \in GL_n$ such that $L(I_n) = BB^T$ and so $H = \Gamma_{B^{-1}} \circ L$ is an isometry of (\mathcal{P}_n, g) fixing I_n .

Let us consider the differential: $D = dH_{I_n} : T_{I_n}(\mathcal{P}_n) \rightarrow T_{I_n}(\mathcal{P}_n)$ (remember that $T_{I_n}(\mathcal{P}_n) = \text{Sym}_n$). We want to prove that $D(I_n) = \pm I_n$.

For, D preserves the metric g and its Riemann tensor of type $(0, 4)$ at I_n . Remembering 2.3 (b), we have $\text{tr}([D(I_n), D(W)]^2) = \text{tr}([I_n, W]^2) = 0$ for every $W \in \text{Sym}_n$. Since the bracket of symmetric matrices is skew-symmetric and since the opposite of the trace of the square of a skew-symmetric matrix is its Frobenius norm, we get that $[D(I_n), D(W)] = 0$ for every $W \in \text{Sym}_n$, i.e. $[D(I_n), U] = 0$ for every $U \in \text{Sym}_n$, because D is bijective and therefore $D(I_n) = \lambda I_n$ for some $\lambda \in \mathbb{R}$. Since D is an isometry, we get $\lambda = \pm 1$.

Now note that the space $(I_n)^\perp := \{W \in \text{Sym}_n : g_{I_n}(I_n, W) = 0\} = \text{Sym}_n^0 = T_{I_n}(SL\mathcal{P}_n)$ is invariant with respect to D , because $D(I_n) = \pm I_n$. Hence D' , the restriction of D to $(I_n)^\perp$, is an isometry of $T_{I_n}(SL\mathcal{P}_n)$ with respect to the metric g .

Since D preserves the Riemann tensor of (\mathcal{P}_n, g) at I_n , D' preserves its restriction to $T_{I_n}(SL\mathcal{P}_n)$, but this last restriction is the Riemann tensor of $(SL\mathcal{P}_n, g)$ at I_n , because $(SL\mathcal{P}_n, g)$ is a totally geodesic submanifold of (\mathcal{P}_n, g) (remember 2.6 (b)).

Since $SL\mathcal{P}_n$ is simply connected, complete and symmetric (remember (3.1)), by

[14, Ch. VI Cor. 7.9], there exists a unique isometry G of $(SL\mathcal{P}_n, g)$ such that $G(I_n) = I_n$ and $dG_{I_n} = D'$.

Now we denote $G^\pm(A, x) = (G(A), \pm x)$ for every $(A, x) \in SL\mathcal{P}_n \times \mathbb{R}$. $G^\pm(A, x)$ are isometries of $(SL\mathcal{P}_n \times \mathbb{R}, g \times h)$ such that $G^\pm(I_n, 0) = (I_n, 0)$ and such that $dG_{(I_n, 0)}^\pm = D' \times (\pm Id_{\mathbb{R}})$.

Easy computations show that $dF_{(I_n, 0)}(V, x) = \frac{x}{\sqrt{n}} I_n + V$ for every $x \in \mathbb{R}$ and every $V \in$

$$T_{I_n}(SL\mathcal{P}_n) = \text{Sym}_n^0 \text{ and that } dF_{I_n}^{-1}(W) = \left(W - \frac{\text{tr}(W)}{n} I_n, \frac{\text{tr}(W)}{\sqrt{n}} \right),$$

for every $W \in T_{I_n}(\mathcal{P}_n) = \text{Sym}_n$, where F and F^{-1} are the mapping recalled above.

Now $(F \circ G^\pm \circ F^{-1})(I_n) = I_n = H(I_n)$ and

$$\begin{aligned} d(F \circ G^\pm \circ F^{-1})_{I_n}(W) &= dF_{(I_n, 0)}(dG_{(I_n, 0)}^\pm(W - \frac{\text{tr}(W)}{n} I_n, \frac{\text{tr}(W)}{\sqrt{n}})) = \\ &= dF_{(I_n, 0)}(dH_{I_n}(W - \frac{\text{tr}(W)}{n} I_n), \pm \frac{\text{tr}(W)}{\sqrt{n}}) = \pm \frac{\text{tr}(W)}{n} I_n + dH_{I_n}(W - \frac{\text{tr}(W)}{n} I_n) = \\ &= \pm \frac{\text{tr}(W)}{n} I_n + dH_{I_n}(W) \mp \frac{\text{tr}(W)}{n} I_n = dH_{I_n}(W) \text{ for every } W \in T_{I_n}(\mathcal{P}_n) = \text{Sym}_n. \end{aligned}$$

Therefore $F \circ G^\pm \circ F^{-1} = H$ (see again [22, Ch. 3 Prop. 62]).

Now $L = \Gamma_B \circ H = \Gamma_B \circ F \circ G^\pm \circ F^{-1}$ and easy computations allow to get the expressions in the statement. \square

Remembering the definitions of φ and ψ in 1.1, from the Theorem 4.1 and 4.4, we

easily get the following

Theorem 4.2. *A mapping $L: (\mathcal{P}_n, g) \rightarrow (\mathcal{P}_n, g)$ is an isometry if and only if there exists a matrix $M \in GL_n$ such that*

$$L = \Gamma_M \text{ or } L = \Gamma_M \circ \varphi \text{ or } L = \Gamma_M \circ \psi \text{ or } L = \Gamma_M \circ \varphi \circ \psi.$$

4.5 Remark. When $n = 2$, we have $\psi = \Gamma_W \circ \varphi = \varphi \circ \Gamma_W$ with W as in 4.3. Hence in the previous Theorem there are only two mutually exclusive possibilities: $L = \Gamma_M$ and $L = \Gamma_M \circ \varphi$.

If $n \geq 3$, since $\varphi, \psi, \varphi \circ \psi$ are not congruences, then the families of isometries, listed in Theorem 4.2, are mutually disjoint. Therefore:

- if $n = 2$ or n is odd, then $I(\mathcal{P}_n, g)$ is a Lie group with four connected components and with $I^0(\mathcal{P}_n, g)$ isomorphic to GL_n^+ if n is odd and $I^0(\mathcal{P}_2, g)$ isomorphic to $GL_2^+ / \{\pm I_2\}$;
- if $n \neq 2$ is even, then $I(\mathcal{P}_n, g)$ is a Lie group with eight connected components and with $I^0(\mathcal{P}_n, g)$ isomorphic to $GL_n^+ / \{\pm I_n\}$.

4.6 Remark. Standard computations allow to obtain the following geometric descriptions of the isometries φ, ψ and $\varphi \circ \psi$ by means of the results of §3 and §4:

- φ is the symmetry with respect to I_n ;
- ψ is the orthogonal symmetry with respect to the hypersurface $SL\mathcal{P}_n$;
- $\varphi \circ \psi$ is the orthogonal symmetry with respect to the geodesic $\mathcal{R} = \{tI_n : t \in \mathbb{R}, t > 0\}$ (i.e. the geodesic through I_n and orthogonal to $SL\mathcal{P}_n$).

4.7 Remark. Let H_n be the real manifold of positive definite hermitian matrices of order n . The tensor g defines also on H_n a structure of Riemannian manifold and (\mathcal{P}_n, g) is one of its Riemannian submanifolds. From [20, Thm. 3] and from the Theorem 4.2 we conclude that every isometry of (\mathcal{P}_n, g) is restriction of an isometry of (H_n, g) .

References

- [1] APOSTOL Tom M., *Calculus*, Vol. II, second edition, John Wiley and Sons, New York, 1969.
- [2] BESSE Arthur L., *Einstein Manifolds*, Springer-Verlag, Berlin, 1987.
- [3] BHATIA Rajendra, *Positive Definite Matrices*, Princeton University Press, Princeton, 2007.
- [4] BHATIA Rajendra, HOLBROOK John, “Riemannian geometry and matrix geometric means”, in *Linear Algebra Appl.* (2006), 413, 594–618.
- [5] BRIDSON Martin R., HAEFLIGER André, *Metric Spaces of Non-Positive Curvature*, GMW 319, Springer-Verlag, Berlin, 1999.
- [6] DOLCETTI Alberto, PERTICI Donato, “Some differential properties of $GL_n(\mathbb{R})$ with the trace metric”, in *Riv. Mat. Univ. Parma* (2015), 6, 267–286.

- [7] DOLCETTI Alberto, PERTICI Donato, “Skew symmetric logarithms and geodesics on $O_n(\mathbb{R})$ ”, *Adv. Geom.*; 18(4) (2018): 495–507.
- [8] FULTON William, HARRIS Joe, *Representation Theory. A First Course*, GTM 129. Springer, New York, 2004.
- [9] HALL Brian C., *Lie Groups, Lie Algebras, and Representations: An Elementary Introduction*, GTM 222, Springer-Verlag, New York, 2003.
- [10] HELGASON Sigurdur, *Differential geometry, Lie groups, and symmetric spaces*, GSM 34, American Mathematical Society, Providence, Rhode Island, 2001.
- [11] HIGHAM Nicholas J., *Functions of Matrices. Theory and Computation*, SIAM Society for Industrial and Applied Mathematics, Philadelphia, 2008.
- [12] HIRSCH Morris W., *Differential Topology*, GTM 33, Springer-Verlag, New York, 1976.
- [13] HORN Roger A., JOHNSON Charles R., *Matrix analysis*, Second edition, Cambridge University Press, Cambridge, 2013.
- [14] KOBAYASHI Shoshichi, NOMIZU Katsumi, *Foundations of differential geometry*, Vol. 1, Interscience Publishers John Wiley and Sons, New York, 1963.
- [15] KOBAYASHI Shoshichi, NOMIZU Katsumi, *Foundations of differential geometry*, Vol. 2, Interscience Publishers John Wiley and Sons, New York, 1969.
- [16] LANG Serge, *Fundamentals of Differential Geometry*, GTM 191, Springer-Verlag, New York, 1999.
- [17] LAWSON Jimmie D., LIM Yongdo, “The Geometric Mean, Matrices, Metrics, and More”, in *The American Mathematical Monthly*, Vol. 108, No.9, (2001), pp.797–812.
- [18] LAWSON H. Blaine Jr, MICHELSON Marie-Louise, *Spin Geometry*, Princeton University Press, Princeton, 1989.
- [19] MOHAKER Maher, ZÉRAÏ Mourad, “The Riemannian Geometry of The Space of Positive-Definite Matrices and Its Application to the Regularization of Positive-Definite Matrix-Valued Data”, in *J. Math. Imaging Vis.*, 40, no. 2, (2011), 171–187.
- [20] MOLNÁR Lajos, “Jordan triple endomorphisms and isometries of spaces of positive definite matrices”, in *Linear and Multilinear Algebra*, 63:1, (2015), 12–33.
- [21] NIELSEN Frank, BHATIA Rajendra (editors), *Matrix Information Geometry*, Springer-Verlag, Berlin, 2013.
- [22] O’NEILL Barrett, *Semi-Riemannian Geometry with applications to relativity*, Academic Press, San Diego, 1983.

- [23] ONISHCHIK Arkadij L., VINBERG Ernest B., “Foundation of Lie Theory”, in ONISHCHIK Arkadij L. (Ed.), *Lie Groups and Lie Algebras I*, Encyclopaedia of Mathematical Sciences Vol. 20, Springer-Verlag, Berlin, 1990.
- [24] SAVAGE Richard P., Jr., “The space of positive definite matrices and Gromov’s invariant”, *Transactions of the American Mathematical Society*, 274 no. 1, (1982), 239–263.
- [25] SKOVGAARD Lene Theil, “A Riemannian Geometry of the Multivariate Normal Model”, *Scandinavian Journal of Statistics*, Vol. 11, No 4, (1984), pp. 211–223.
- [26] VAN LEEUVEN Mark A. A., COHEN Arjeh M., LISSER Bert, *LiE, A Package for Lie Group Computations*, Computer Algebra Nederland, Amsterdam, 1992. Manual and on-line service available on <http://www.mathlabo.univ-poitiers.fr/~maavl/LiE/>, accessed April 24th, 2018.
- [27] WARNER Frank W., *Foundations of differentiable manifolds and Lie groups*, Springer-Verlag, New York, 1983.
- [28] WOLF Joseph A., *Spaces of Constant Curvature*, Sixth Edition, AMS Chelsea Publishing, Providence, 2011.

Mathematics Subject Classification (2010): 15B48, 53C35, 53C50, 17B40, 17B10

Università di Firenze, Dipartimento di Matematica e Informatica “Ulisse Dini”, Viale Morgagni 67/a, 50134 Firenze, ITALIA
e-mail: alberto.dolcetti@unifi.it,
<http://orcid.org/0000-0001-9791-8122>
e-mail: donato.pertici@unifi.it,
<http://orcid.org/0000-0003-4667-9568>

Lavoro pervenuto in redazione il MM.GG.AAAA.